

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|--|--|--|
| (51) International Patent Classification ⁶ : C07D 487/04, A61K 31/495 | A1 | (11) International Publication Number: WO 97/29109 (43) International Publication Date: 14 August 1997 (14.08.97) |
| (21) International Application Number: PCT/EP97/00459 (22) International Filing Date: 30 January 1997 (30.01.97) (30) Priority Data: 60/011,279 7 February 1996 (07.02.96) US 60/027,688 8 October 1996 (08.10.96) US (71) Applicants (for all designated States except US): JANSSEN PHARMACEUTICA N.V. [BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE). NEUROCRINE BIOSCIENCES INC. [US/US]; 3050 Science Park Road, San Diego, CA 92121-1102 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): CHEN, Chen [CN/US]; Neurocrine Biosciences Inc., 3050 Science Park Road, San Diego, CA 92121-1102 (US). WEBB, Thomas, R. [US/US]; Neurocrine Biosciences Inc., 3050 Science Park Road, San Diego, CA 92121-1102 (US). McCARTHY, James, R. [US/US]; Neurocrine Biosciences Inc., 3050 Science Park Road, San Diego, CA 92121-1102 (US). MORAN, Terence, J. [US/US]; Neurocrine Biosciences Inc., 3050 Science Park Road, San Diego, CA 92121-1102 (US). WILCOXEN, Keith, M. [US/US]; Neurocrine Biosciences Inc., 3050 Science Park Road, San Diego, CA 92121-1102 (US). | (74) Agent: WANTE, Dirk; Janssen Pharmaceutica N.V., Patent Dept., Turnhoutseweg 30, B-2340 Beerse (BE). (81) Designated States: AL, AM, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KG, KR, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report. | |

(54) Title: PYRAZOLOPYRIMIDINES AS CRF RECEPTOR ANTAGONISTS

(57) Abstract

This invention concerns compounds of formula (I), including the stereoisomers and the pharmaceutically acceptable acid addition salt forms thereof, wherein R¹ is NR⁴R⁵ or OR⁵; R² is C₁-alkyl, C₁-alkyloxy or C₁-alkylthio; R³ is hydrogen, C₁-alkyl, C₁-alkylsulfonyl, C₁-alkylsulfoxy or C₁-alkylthio; R⁴ is hydrogen, C₁-alkyl, mono- or di(C₃-cycloalkyl)methyl, C₃-cycloalkyl, C₃-alkenyl, hydroxyc₁-alkyl, C₁-alkylcarbonyloxyC₁-alkyl or C₁-alkyloxyC₁-alkyl; R⁵ is C₁-alkyl, mono- or di(C₃-cycloalkyl)methyl, Ar¹CH₂, C₁-alkyloxyC₁-alkyl, hydroxyc₁-alkyl, C₃-alkenyl, thienylmethyl, furanymethyl, C₁-alkylthioC₁-alkyl, morpholinyl, mono- or di(C₁-alkyl)aminoC₁-alkyl, di(C₁-alkyl)amino, C₁-alkylcarbonylC₁-alkyl, C₁-alkyl substituted with imidazolyl; or a radical of the formula -Alk-O-CO-Ar¹; or R⁴ and R⁵ taken together with the nitrogen atom to which they are attached may form an optionally substituted pyrrolidinyl, piperidinyl, homopiperidinyl or morpholinyl group; having CRF receptor antagonistic properties; pharmaceutical compositions containing such compounds as active ingredients; methods of treating disorders related to hypersecretion of CRF such as depression, anxiety, substance abuse, by administering an effective amount of a compound of formula (I).



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | |
|----|--------------------------|----|--|----|--------------------------|
| AM | Armenia | GB | United Kingdom | MW | Malawi |
| AT | Austria | GE | Georgia | MX | Mexico |
| AU | Australia | GN | Guinea | NE | Niger |
| BB | Barbados | GR | Greece | NL | Netherlands |
| BE | Belgium | HU | Hungary | NO | Norway |
| BF | Burkina Faso | IE | Ireland | NZ | New Zealand |
| BG | Bulgaria | IT | Italy | PL | Poland |
| BJ | Benin | JP | Japan | PT | Portugal |
| BR | Brazil | KE | Kenya | RO | Romania |
| BY | Belarus | KG | Kyrgyzstan | RU | Russian Federation |
| CA | Canada | KP | Democratic People's Republic of Korea | SD | Sudan |
| CF | Central African Republic | KR | Republic of Korea | SE | Sweden |
| CG | Congo | KZ | Kazakhstan | SG | Singapore |
| CH | Switzerland | LI | Liechtenstein | SI | Slovenia |
| CI | Côte d'Ivoire | LK | Sri Lanka | SK | Slovakia |
| CM | Cameroon | LR | Liberia | SN | Senegal |
| CN | China | LT | Lithuania | SZ | Swaziland |
| CS | Czechoslovakia | LU | Luxembourg | TD | Chad |
| CZ | Czech Republic | LV | Latvia | TG | Togo |
| DE | Germany | MC | Monaco | TJ | Tajikistan |
| DK | Denmark | MD | Republic of Moldova | TT | Trinidad and Tobago |
| EE | Estonia | MG | Madagascar | UA | Ukraine |
| ES | Spain | ML | Mali | UG | Uganda |
| FI | Finland | MN | Mongolia | US | United States of America |
| FR | France | MR | Mauritania | UZ | Uzbekistan |
| GA | Gabon | | | VN | Viet Nam |

PYRAZOLOPYRIMIDINES AS CRF RECEPTOR ANTAGONISTS

Background of the invention

- 5 This invention relates to pyrazolopyrimidines which possess CRF receptor antagonistic properties, to pharmaceutical compositions containing these compounds as active ingredient, and the use thereof in the treatment of endocrine, psychiatric and neurologic conditions or illnesses, including stress-related disorders in general.
- 10 The first corticotropin-releasing factor (CRF) was isolated from ovine hypothalmi and identified as a 41-amino acid peptide (Vale et al., *Science* 213:1394-1397, 1981). Subsequently, sequences of human and rat CRF were isolated and determined to be identical, but different from ovine CRF in 7 of the 41 amino acid residues (Rivier et al., *Proc. Natl. Acad. Sci. USA* 80:4851, 1983; Shibahara et al., *EMBO J.* 2:775, 1983).
- 15 CRF has been found to produce profound alterations in endocrine, nervous and immune system functions. CRF is believed to be the major physiological regulator of the basal and stress-release of adrenocorticotrophic hormone ("ACTH"), β -endorphin, and other pro-opiomelanocortin ("POMC")-derived peptides from the anterior pituitary (Vale et al., *Science* 213:1394-1397, 1981). Briefly, CRF is believed to initiate its biological
- 20 effects by binding to a plasma membrane receptor which has been found to be distributed throughout the brain (DeSouza et al., *Science* 221:1449-1451, 1984), pituitary (DeSouza et al., *Methods Enzymol.* 124:560, 1986; Wynn et al., *Biochem. Biophys. Res. Comm.* 110:602-608, 1983), adrenals (Udelsman et al., *Nature* 319:147-150, 1986) and spleen (Webster, E.L., and E.B. DeSouza, *Endocrinology* 122:609-617, 1988). The CRF receptor is coupled to a GTP-binding protein (Perrin et al., *Endocrinology* 118: 1171- 1179, 1986) which mediates CRF-stimulated increase in intracellular production of cAMP (Bilezikjian, L.M., and W.W. Vale, *Endocrinology* 113:657-662, 1983).
- 30 In addition to its role in stimulating the production of ACTH and POMC, CRF is also believed to coordinate many of the endocrine autonomic, and behavioral responses to stress, and may be involved in the pathophysiology of affective disorders. Moreover, CRF is believed to be a key intermediary in communication between the immune, central nervous, endocrine and cardiovascular systems (Crofford et al., *J. Clin. Invest.* 90:2555-2564, 1992; Sapolsky et al., *Science* 238:522-524, 1987; Tilders et al., *Regul. Peptides* 5:77-84, 1982). Overall, CRF appears to be one of the pivotal central nervous
- 35 system neurotransmitters and plays a crucial role in integrating the body's overall response to stress.

Administration of CRF directly to the brain elicits behavioral, physiological, and endocrine responses identical to those observed for an animal exposed to a stressful environment. For example, intracerebroventricular injection of CRF results in behavioral activation (Sutton et al., *Nature* 297:331, 1982), persistent activation of the electroencephalogram (Ehlers et al., *Brain Res.* 2/8332, 1983), stimulation of the sympathoadrenomedullary pathway (Brown et al., *Endocrinology* 110:928, 1982), an increase of heart rate and blood pressure (Fisher et al., *Endocrinology* 110:2222, 1982), an increase in oxygen consumption (Brown et al., *Life Sciences* 30:207, 1982), alteration of gastrointestinal activity (Williams et al., *Am. J. Physiol.* 253:G582, 1987), suppression of food consumption (Levine et al., *Neuropharmacology* 22:337, 1983), modification of sexual behavior (Sirinathsinghji et al., *Nature* 305:232, 1983), and immune function compromise (Irwin et al., *Am. J. Physiol.* 255:R744, 1988). Furthermore, clinical data suggest that CRF may be hypersecreted in the brain in depression, anxiety-related disorders, and anorexia nervosa. (DeSouza, *Ann. Reports in Med. Chem.* 25:215-223, 1990).

Accordingly, clinical data suggest that CRF receptor antagonists may represent novel antidepressant and/or anxiolytic drugs that may be useful in the treatment of the neuropsychiatric disorders manifesting hypersecretion of CRF.


CRF receptor antagonists have been reported in for example, U.S. Patent No. 5,063,245 disclosing substituted 4-thio-5-oxo-3-pyrazoline derivatives and Australian Patent No. AU-A-41399/93, disclosing substituted 2-aminothiazole derivatives. WO-92/18504 and JP-32/04877 disclose pyrazolo[1,5-a]pyrimidines as antiinflammatory agents. Also, WO-94/13676, WO-94/13677 and WO-95/33750 disclose pyrrolopyrimidines, pyrazolo[3,4-d]pyrimidines and substituted purines as CRF receptor antagonists.

Arylpyrazolo[1,5-a]pyrimidines have been described as xanthine oxidase inhibitors (Robins et al., *J. Heterocyclic Chem.* 22:601-634, 1985). JP-42/011,753 discloses 7-methylamino-pyrazolo[1,5-a]pyrimidine derivatives useful as sedative and antiphlogistic agents. And JP-61/057,587 discloses pyrazolo[1,5-a]pyrimidine derivatives useful as antiulcer agents.

Due to the physiological significance of CRF, the development of further biologically active small molecules having significant CRF receptor binding activity and which are capable of antagonizing the CRF receptor remains a desirable goal. Such CRF receptor antagonists would be useful in the treatment of endocrine, psychiatric and neurologic conditions or illnesses, including stress-related disorders in general.

Description of the invention

This invention concerns CRF antagonistic compounds of formula (I)



5

including the stereoisomers and the pharmaceutically acceptable acid addition salt forms thereof, wherein

R¹ is NR⁴R⁵ or OR⁵;

R² is C₁₋₆alkyl, C₁₋₆alkyloxy or C₁₋₆alkylthio;

10 R³ is hydrogen, C₁₋₆alkyl, C₁₋₆alkylsulfonyl, C₁₋₆alkylsulfoxy or C₁₋₆alkylthio;

R⁴ is hydrogen, C₁₋₆alkyl, mono- or di(C₃₋₆cycloalkyl)methyl, C₃₋₆cycloalkyl, C₃₋₆alkenyl, hydroxyC₁₋₆alkyl, C₁₋₆alkylcarbonyloxyC₁₋₆alkyl or C₁₋₆alkyloxyC₁₋₆alkyl;

15 R⁵ is C₁₋₈alkyl, mono- or di(C₃₋₆cycloalkyl)methyl, Ar¹CH₂, C₁₋₆alkyloxyC₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₃₋₆alkenyl, thienylmethyl, furanymethyl, C₁₋₆alkylthioC₁₋₆alkyl, morpholinyl, mono- or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, di(C₁₋₆alkyl)amino, C₁₋₆alkylcarbonylC₁₋₆alkyl, C₁₋₆alkyl substituted with imidazolyl; or a radical of formula -Alk-O-CO-Ar¹;

20 or R⁴ and R⁵ taken together with the nitrogen atom to which they are attached may form a pyrrolidinyl, piperidinyl, homopiperidinyl or morpholinyl group, optionally substituted with C₁₋₆alkyl or C₁₋₆alkyloxyC₁₋₆alkyl; and

25 Ar is phenyl; phenyl substituted with 1, 2 or 3 substituents independently selected from halo, C₁₋₆alkyl, trifluoromethyl, hydroxy, cyano, C₁₋₆alkyloxy, benzyloxy, C₁₋₆alkylthio, nitro, amino and mono- or di(C₁₋₆alkyl)amino; pyridinyl; pyridinyl substituted with 1, 2 or 3 substituents independently selected from halo, C₁₋₆alkyl, trifluoromethyl, hydroxy, cyano, C₁₋₆alkyloxy, benzyloxy, C₁₋₆alkylthio, nitro, amino, mono- or di(C₁₋₆alkyl)amino and piperidinyl; and wherein said substituted phenyl may optionally be further substituted with one or more halogens;

30 Ar¹ is phenyl; phenyl substituted with 1, 2 or 3 substituents each independently selected from halo, C₁₋₆alkyl, C₁₋₆alkyloxy, di(C₁₋₆alkyl)aminoC₁₋₆alkyl, trifluoromethyl and C₁₋₆alkyl substituted with morpholinyl; or pyridinyl; and

Alk is C₁₋₆alkanediyl.

In a further aspect the invention concerns novel compounds of formula (I) as defined above, with the proviso that 5-methyl-3-phenyl-7-(phenylmethoxy)-pyrazolo[1,5-a]-pyrimidine and 2,5-dimethyl-7-(methylamino)-3-phenyl-pyrazolo[1,5-a]pyrimidine are not included

The proviso is intended to exclude compounds disclosed in JP-61/057,587 and JP-42/011,753.

As used in the foregoing definitions and hereinafter, halo is generic to fluoro, chloro, bromo and iodo; C₁₋₆alkanediyl defines bivalent straight and branched chained saturated hydrocarbon radicals having from 1 to 6 carbon atoms, such as, for example, methylene, 1,2-ethanediyl, 1,3-propanediyl, 1,4-butanediyl, 1,5-pentanediyl, 1,6-hexanediyl and the branched isomers thereof; C₁₋₂alkyl defines straight saturated hydrocarbon radicals having from 1 to 2 carbon atoms such as methyl and ethyl; C₂₋₄alkyl defines straight and branched chain saturated hydrocarbon radicals having from 2 to 4 carbon atoms such as ethyl, propyl, butyl, 1-methylethyl and the like; C₃₋₄alkyl defines straight and branched chain saturated hydrocarbon radicals having from 3 to 4 carbon atoms such as propyl, butyl, 1-methylethyl and the like; C₁₋₆alkyl includes C₁₋₂alkyl and C₃₋₄alkyl radicals as defined hereinbefore and the higher homologs thereof having from 5 to 6 carbon atoms such as, pentyl, the pentyl isomers, hexyl and the hexyl isomers; C₁₋₈alkyl includes C₁₋₆alkyl and the higher homologues thereof having from 7 to 8 carbon atoms such as, for example, heptyl, octyl and the like; C₃₋₆alkenyl defines straight and branched chain hydrocarbon radicals containing one double bond and having from 3 to 6 carbon atoms such as, for example, 2-propenyl, 3-butenyl, 2-pentenyl, 3-pentenyl, 3-methyl-2-butenyl, and the like; and where said C₃₋₆alkenyl is linked to a nitrogen or oxygen, the carbon atom making the link preferably is saturated. C₃₋₆cycloalkyl comprises cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. HydroxyC₁₋₆alkyl refers to C₁₋₆alkyl substituted with a hydroxyl-group. Homopiperidinyl refers to a 7 membered saturated ring containing one nitrogen atom.

Depending on the nature of some of the substituents, the compounds of formula (I) may contain one or more asymmetric centers which may be designated with the generally used R and S nomenclature.

- The compounds of the present invention contain basic nitrogen atoms and, as such, can be present as the free base or in the form of acid addition salts, both being part of this invention. Acid addition salts may be prepared by methods well known in the art, and may be formed from organic and inorganic acids. Suitable organic acids include maleic, fumaric, benzoic, ascorbic, succinic, methanesulfonic, acetic, oxalic, propionic, tartaric, salicylic, citric, gluconic, lactic, mandelic, cinnamic, aspartic, stearic, palmitic, glycolic, glutamic, and benzenesulfonic acids. Suitable inorganic acids include hydrochloric, hydrobromic, sulfuric, phosphoric, and nitric acids.
- 10 Particular groups of compounds within the invention are those compounds of formula (I) wherein one or more of the following restrictions apply :
- a) R^1 is NR^4R^5 wherein R^4 is hydrogen, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, C_{1-6} alkylcarbonyloxy C_{1-6} alkyl or C_{3-6} alkenyl; in particular C_{2-4} alkyl, hydroxy C_{1-2} alkyl, C_{3-4} alkenyl or C_{1-2} alkylcarbonyloxy C_{2-4} alkyl; and R^5 is C_{1-8} alkyl, C_{3-6} alkenyl, hydroxy C_{1-6} alkyl, C_{1-6} alkyloxy C_{1-6} alkyl, phenylmethyl or C_{3-6} cycloalkylmethyl; in particular C_{2-4} alkyl, C_{3-4} alkenyl, hydroxy C_{2-4} alkyl or cyclopropylmethyl;
 - b) R^1 is OR^5 wherein R^5 is C_{1-6} alkyl; in particular C_{2-4} alkyl;
 - c) R^2 is C_{1-6} alkyl; in particular C_{1-2} alkyl;
 - 20 d) R^3 is hydrogen, C_{1-6} alkyl or C_{1-6} alkylthio; in particular hydrogen, C_{1-2} alkyl or C_{1-2} alkylthio;
 - e) Ar is a phenyl substituted with 1, 2 or 3 substituents each independently selected from C_{1-6} alkyl, C_{1-6} alkyloxy or halo; wherein the phenyl moiety is preferably substituted in the 3-, 4-, 6-, 2,4- or 2,4,6-positions; or Ar is a pyridinyl substituted with 1, 2 or 3 substituents each independently selected from halo, amino, nitro, trifluoromethyl, mono- or di(C_{1-6} alkyl)amino, piperidinyl or C_{1-6} alkyl; wherein the pyridinyl moiety preferably is connected via the 2- or 3-position to the remainder of the molecule.
- 30 Another particular group of compounds are those compounds of formula (I) wherein R^1 is NR^4R^5 and R^4 and R^5 are taken together with the nitrogen atom to which they are attached to form a pyrrolidinyl, piperidinyl, homopiperidinyl or morpholinyl group, optionally substituted with C_{1-6} alkyl or C_{1-6} alkyloxy C_{1-6} alkyl.
- 35 Preferred compounds are those compounds of formula (I) wherein R^1 is NR^4R^5 and R^4 is C_{3-4} alkyl or allyl, preferably propyl; R^5 is C_{2-4} alkyl, allyl or cyclopropylmethyl, preferably propyl; R^2 is methyl; R^3 is hydrogen, methyl or methylthio, preferably

propyl; and Ar is a phenyl substituted with 1, 2 or 3 substituents each independently selected from halo, methyl or methoxy; and Ar in particular is pyridinyl substituted with 1, 2 or 3 substituents each independently selected from halo, methyl or dimethylamino.

5

More preferably Ar is 3-pyridinyl substituted in the 4- and/or 6-position with methyl or dimethylamino.

Most preferred are those compounds selected from

10

3-(2,4-dichlorophenyl)-5-methyl-7-(*N*-propyl-*N*-cyclopropanemethylamino)-pyrazolo[2,3-*a*]pyrimidine;

3-(2,4-dichlorophenyl)-5-methyl-7-(*N*-allyl-*N*-cyclopropanemethylamino)-pyrazolo[2,3-*a*]pyrimidine;

15

2-methylthio-3-(2,4-dichlorophenyl)-5-methyl-7-(*N,N*-diallylamino)-pyrazolo[2,3-*a*]pyrimidine;

2-methylthio-3-(2,4-dichlorophenyl)-5-methyl-7-(*N*-butyl-*N*-cyclopropanemethylamino)pyrazolo[2,3-*a*]pyrimidine;

2-methylthio-3-(2,4-dichlorophenyl)-5-methyl-7-(*N*-propyl-*N*-cyclopropanemethylamino)pyrazolo[2,3-*a*]pyrimidine;

20

2-methyl-3-(4-chlorophenyl)-5-methyl-7-(*N,N*-dipropylamino)-pyrazolo[2,3-*a*]pyrimidine;

3-[6-(dimethylamino)-3-pyridinyl]-2,5-dimethyl-*N,N*-dipropylpyrazolo[2,3-*a*]pyrimidin-7-amine; or

3-[6-(dimethylamino)-4-methyl-3-pyridinyl]-2,5-dimethyl-*N,N*-dipropylpyrazolo[2,3-*a*]pyrimidine-7-amine; or

25

3-(2,4-dimethoxyphenyl)-2,5-dimethyl-7-(*N*-propyl-*N*-methoxyethylamino)-pyrazolo[2,3-*a*]pyrimidine;


the stereoisomeric forms and the pharmaceutically acceptable acid addition salts thereof.

30

The compounds of the present invention can generally be prepared by alkylating a pyrazolopyrimidine of formula (II) with an intermediate of formula (III). In intermediate (II), W is an appropriate leaving group such as halo, e.g. chloro, bromo, or a sulfonyloxy group, e.g. a mesyloxy or a tosyloxy group.


35

-7-




The above reaction is typically conducted in a suitable solvent, e.g. an aprotic solvent such as DMF or acetonitrile, an ether, e.g. tetrahydrofuran, preferably at an elevated temperature and, when the intermediates of formula (III) are volatile amines, in a sealed reaction vial.

Also, compounds of formula (I) wherein R^1 is OR^5 , said compounds being represented by formula (I-a), may be prepared by *O*-alkylating an intermediate of formula (VI) with an intermediate of formula (VII), wherein W is as defined above.



Said reaction for preparing compounds of formula (I-a) can be performed in a reaction-inert solvent such as, for example, *N,N*-dimethylformamide, and in the presence of a suitable base such as, for example, sodium hydride, preferably at a temperature ranging between room temperature and reflux temperature.

The compounds of formula (I) wherein R^1 is NR^4R^5 , represented by formula (I-c), can be prepared from either compounds of formula (VIII) or (IX) by suitable *N*-alkylation reactions as depicted herebelow, wherein W is as previously defined. These *N*-alkylations are conducted in a reaction-inert solvent such as, for example, an ether e.g. tetrahydrofuran and preferably in the presence of a strong base, e.g. NaH.



In certain instances, this reaction can give rise to side products wherein R² is alkylated by (R⁴ or R⁵)-W, in particular where R² is methyl and R⁴ or R⁵ is C₁₋₆alkyl.

5 As outlined below, compounds of formula (I) may be converted into each other following art-known transformation procedures.


10 The compounds of formula (I) wherein R³ is C₁₋₆alkylthio can be converted into compounds of formula (I) wherein R³ is C₁₋₆alkylsulfonyl or C₁₋₆alkylsulfoxy by an oxidation reaction, e.g. treatment with a peroxide such as 3-chloroperbenzoic acid in a reaction-inert solvent, e.g. dichloromethane. By controlling the amount of oxidant and other reaction parameters, either compounds of formula (I) wherein R³ is C₁₋₆alkylsulfonyl or C₁₋₆alkylsulfoxy can be obtained, or a mixture of both, which subsequently can be separated by conventional methods, e.g. column chromatography.

15 The compounds of formula (I) may also be converted into each other via art-known reactions or functional group transformations. For instance, compounds of formula (I) bearing a hydroxyC₁₋₆alkyl group may be converted into compounds of formula (I) bearing a C₁₋₆alkylcarbonyloxyC₁₋₆alkyl group, e.g. by treatment with an acid anhydride in a reaction-inert solvent such as, e.g. dichloromethane, and optionally in the presence of a base such as, e.g. pyridine.

20 Compounds of formula (I) bearing a nitro group may be converted to compounds of formula (I) bearing an amino group and subsequently to compounds of formula (I) having a mono- or di(C₁₋₆alkyl)amino group. Also, the amino group may be converted using a diazotization reaction to a halo.


30 Further, the Ar group of compounds of formula (I) can be halogenated using a halogenating agent such as, e.g. chlorine or bromine, in a suitable solvent, e.g. acetic acid, and optionally the reaction may be performed at a temperature ranging between room temperature and the reflux temperature of the reaction mixture.

Intermediates of formula (II) can be prepared according to the procedure as described in Robins et al., *J. Heterocyclic Chem.* 22:601-634, 1985.



- Aminopyrazole derivatives (IV) are reacted with a β -keto ester (V), preferably under reflux conditions and in a suitable reaction-inert solvent such as an ether, e.g. THF, yielding hydroxypyrazolopyrimidines (VI) which are converted into intermediates of formula (II) by converting the hydroxy group of intermediate (VI) into leaving group W, e.g. by treating (VI) with methanesulfonyloxy chloride or a halogenating reagent such as, e.g. POCl_3 .
- Intermediates of formula (VIII) are prepared by treating intermediates of formula (II) with ammonia.

- In an embodiment, this invention also provides for intermediates of formula (II'), wherein W' represents hydroxy, halo, mesyloxy or tosyloxy; with the proviso that Ar is other than phenyl.



- Said intermediates of formula (II') may be prepared according to procedures used to prepare intermediates of formula (II), thereby thereby yielding compounds of formula (II'-a), defined as compounds of formula (II') wherein W' is hydroxy; and optionally converting compounds of formula (II'-a) into compounds of formula (II'-b), defined as compounds of formula (II') wherein W' is other than hydroxy.
- Stereoisomers may be prepared by separation of the end products of formula (I) following art-known procedures, e.g. by treatment with an optically active acid and separating the thus-formed diastereoisomeric salts by selective crystallization or column chromatography. Or, stereoisomers may be prepared by using stereoisomeric starting materials in any of the above reaction schemes or in the preparation of intermediates described hereinafter.

The effectiveness of a compound as a CRF receptor antagonist may be determined by various assay methods. Suitable CRF antagonists of this invention are capable of inhibiting the specific binding of CRF to its receptor and antagonizing activities associated with CRF. A compound of structure (I) may be assessed for activity as a CRF antagonist by one or more generally accepted assays for this purpose, including (but not limited to) the assays disclosed by DeSouza et al. (*J. Neuroscience* 7:88, 1987) and Battaglia et al. (*Synapse* 1:572, 1987). As mentioned above, suitable CRF antagonists include compounds which demonstrate CRF receptor affinity. CRF receptor affinity may be determined by binding studies that measure the ability of a compound to inhibit the binding of a radiolabeled CRF (e.g. [¹²⁵I]tyrosine CFR) to receptor (e.g., receptors prepared from rat cerebral cortex membranes). The radioligand binding assay described by DeSouza et al. (*supra*, 1987) provides an assay for determining a compound's affinity for the CRF receptor. Such activity is typically calculated from the IC₅₀ as the concentration of a compound necessary to displace 50% of the radiolabeled ligand from the receptor, and is reported as a "K_i" value calculated by the following equation :

$$K_i = \frac{IC_{50}}{1 + L / K_D}$$

where L = radioligand and K_D = affinity of radioligand for receptor (Cheng and Prusoff, *Biochem. Pharmacol.* 22:3099, 1973).

In addition to inhibiting CRF receptor binding, a compound's CRF receptor antagonist activity may be established by the ability of the compound to antagonize an activity associated with CRF. For example, CRF is known to stimulate various biochemical processes, including adenylate cyclase activity. Therefore, compounds may be evaluated as CRF antagonists by their ability to antagonize CRF-stimulated adenylate cyclase activity by, for example, measuring cAMP levels. The CRF-stimulated adenylate cyclase activity assay described by Battaglia et al. (*supra*, 1987) provides an assay for determining a compound's ability to antagonize CRF activity. Accordingly, CRF receptor antagonist activity may be determined by assay techniques which generally include an initial binding assay (such as disclosed by DeSouza (*supra*, 1987)) followed by a cAMP screening protocol (such as disclosed by Battaglia (*supra*, 1987)). With reference to CRF receptor binding affinities, CRF receptor antagonists of this invention have a K_i of less than 10 μM. In a preferred embodiment of this invention, a

CRF receptor antagonist has a K_i of less than 1 μM , and more preferably less than 0.25 μM (*i.e.*, 250 nM).

The CRF receptor antagonists of the present invention demonstrate activity at the CRF receptor site, and may be used as therapeutic agents for the treatment of a wide range of disorders or illnesses including endocrine, psychiatric, and neurologic disorders or illnesses. More specifically, the CRF receptor antagonists of the present invention may be useful in treating physiological conditions or disorders arising from the hypersecretion of CRF. Because CRF is believed to be a pivotal neurotransmitter that activates and coordinates the endocrine, behavioral and automatic responses to stress, the CRF receptor antagonists of the present invention can be used to treat neuropsychiatric disorders. Neuropsychiatric disorders which may be treatable by the CRF receptor antagonists of this invention include affective disorders such as depression; anxiety-related disorders such as generalized anxiety disorder, panic disorder, obsessive-compulsive disorder, abnormal aggression, cardiovascular abnormalities such as unstable angina and reactive hypertension; and feeding disorders such as anorexia nervosa, bulimia, and irritable bowel syndrome. CRF antagonists may also be useful in treating stress-induced immune suppression associated with various diseases states, as well as stroke. Other uses of the CRF antagonists of this invention include treatment of inflammatory conditions (such as rheumatoid arthritis, uveitis, asthma, inflammatory bowel disease and G.I. motility), Cushing's disease, infantile spasms, epilepsy and other seizures in both infants and adults, and various substance abuse and withdrawal (including alcoholism).

In another embodiment of the invention, pharmaceutical compositions containing one or more CRF receptor antagonists are disclosed. For the purposes of administration, the compounds of the present invention may be formulated as pharmaceutical compositions. Pharmaceutical compositions of the present invention comprise a CRF receptor antagonist of the present invention (*i.e.*, a compound of structure (I)) and a pharmaceutically acceptable carrier and/or diluent. The CRF receptor antagonist is present in the composition in an amount which is effective to treat a particular disorder, that is, in an amount sufficient to achieve CRF receptor antagonist activity, and preferably with acceptable toxicity to the patient. Preferably, the pharmaceutical compositions of the present invention may include a CRF receptor antagonist in an amount from 0.1 mg to 250 mg per dosage depending upon the route of administration, and more preferably from 1 mg to 60 mg. Appropriate concentrations and dosages can be readily determined by one skilled in the art.

Pharmaceutically acceptable carrier and/or diluents are familiar to those skilled in the art. For compositions formulated as liquid solutions, acceptable carriers and/or diluents include saline and sterile water, and may optionally include antioxidants, buffers, bacteriostats and other common additives. The compositions can also be formulated as pills, capsules, granules, or tablets which contain, in addition to a CRF receptor antagonist, diluents, dispersing and surface active agents, binders, and lubricants. One skilled in this art may further formulate the CRF receptor antagonist in an appropriate manner, and in accordance with accepted practices, such as those disclosed in *Remington's Pharmaceutical Sciences*, Gennaro, Ed., Mack Publishing Co., Easton, USA, 1990.

In another embodiment, the present invention provides a method for treating a variety of disorders or illnesses, including endocrine, psychiatric and neurologic disorders or illnesses. Such methods include administering of a compound of the present invention to a warm-blooded animal in an amount sufficient to treat the disorder or illness. Such methods include systemic administration of a CRF receptor antagonist of this invention, preferably in the form of a pharmaceutical composition. As used herein, systemic administration includes oral and parenteral methods of administration. For oral administration, suitable pharmaceutical compositions of CRF receptor antagonists include powders, granules, pills, tablets, and capsules as well as liquids, syrups, suspensions, and emulsions. These compositions may also include flavorants, preservatives, suspending, thickening and emulsifying agents, and other pharmaceutically acceptable additives. For parental administration, the compounds of the present invention can be prepared in aqueous injection solutions which may contain, in addition to the CRF receptor antagonist, buffers, antioxidants, bacteriostats, and other additives commonly employed in such solutions.

As mentioned above, administration of a compound of the present invention can be used to treat a wide variety of disorders or illnesses. In particular, the compounds of the present invention may be administered to a warm-blooded animal for the treatment of depression, anxiety disorder, panic disorder, obsessive-compulsive disorder, abnormal aggression, unstable angina, reactive hypertension, anorexia nervosa, bulimia, irritable bowel syndrome, stress-induced immune suppression, stroke, inflammation, Cushing's disease, infantile spasms, epilepsy, and substance abuse or withdrawal.

Hence, this invention provides the use of compounds of formula (I) for the manufacture of a medicine for treating physiological conditions or disorders arising from the hypersecretion of corticotropin-releasing factor (CRF); and in a further embodiment the use of novel compounds of formula (I) as a medicine is provided.

The following examples are provided for purposes of illustration, not limitation.

Experimental part

Hereinafter "THF" means tetrahydrofuran and "DCM" means dichloromethane.

A. Preparation of the intermediates.

Example A.1

a) 3-Amino-4-(2,4-dichlorophenyl)pyrazole and ethyl acetoacetate (2 equivalents) were dissolved in dioxane and heated under reflux overnight. The mixture was concentrated *in vacuo* and diluted with ethyl acetate. An off-white solid formed after 2 days standing was collected by vacuum filtration, yielding 3-(2,4-dichlorophenyl)-5-methyl-7-hydroxypyrazolo[2,3-a]pyrimidine (intermediate 1).

b) Intermediate 1 (300 mg) was mixed with POCl_3 (1.5 ml) and heated to reflux for 1 hour. The resultant purple solution was carefully transferred into ice-water. The product was extracted with ethyl acetate, washed with saturated sodium bicarbonate and brine, dried over MgSO_4 and concentrated *in vacuo* to give 3-(2,4-dichlorophenyl)-5-methyl-7-chloropyrazolo[2,3-a]pyrimidine (intermediate 2) as a brown solid (260 mg, 82%).

Example A.2

a) To a stirred solution of sodium hydride (60%, 25 mmol) in THF (10 ml) 6-(dimethylamino)-3-pyridineacetonitrile was added dropwise (10 mmol) in THF (10 ml). The solution was allowed to stir for 10 minutes before ethyl acetate (30 mmol) was added slowly. The resulting suspension was stirred at room temperature for another hour. The reaction mixture was concentrated under vacuum and dissolved in ethyl acetate/methanol (1:1) and filtered through silica. The filtrate was concentrated, yielding (intermediate 7).

b) A mixture of intermediate 7 and hydrazine hydrobromide (100 mmol) was dissolved in ethanol/water (9:1, total 100 ml) and refluxed for 1 hour. The reaction mixture was concentrated and the residue was partitioned between ethyl acetate and sodium bicarbonate solution. The combined extracts were dried over sodium sulfate, filtered and concentrated to dryness. The residue was dissolved in 1,4-dioxane (200 ml) and

refluxed for 16 hours in the presence of ethyl acetoacetate. The reaction mixture was cooled and an off-white solid precipitated out. Diethyl ether was added to aid in crystallization and the precipitate was filtered off and dried, yielding 3-[2-(dimethylamino)-5-pyridinyl]-2,5-dimethyl-7-hydroxypyrazolo[2,3-a]pyrimidine

5 (intermediate 8).


c) Intermediate 8 was dissolved in POCl_3 (2 ml) and refluxed for 2 hours. The reaction mixture was cooled and poured onto ice. The solution was made basic ($\text{pH} \approx 9$) by addition of solid sodium carbonate and extracted with diethyl ether. The combined organic layers were dried over sodium sulfate, filtered and concentrated, yielding 3-[6-(dimethylamino)-3-pyridinyl]-2,5-dimethyl-7-chloro-pyrazolo[2,3-a]pyrimidine

10 (intermediate 9).

Table 1 lists the intermediates that were prepared according to one of the above Examples.

15

Table 1 :



| Interm. No. | Ex. No. | R ³ | Ar |
|-------------|---------|---------------------------------|--------------------------------------|
| 2 | A.1 | H | 2,4-dichlorophenyl |
| 3 | A.1 | CH ₃ S | 2,4-dichlorophenyl |
| 4 | A.1 | CH ₃ | 2,4-dichlorophenyl |
| 5 | A.1 | H | 4-chlorophenyl |
| 6 | A.1 | H | 2,6-dichlorophenyl |
| 13 | A.1 | CH ₃ | 4-chlorophenyl |
| 14 | A.1 | CH ₃ | 3-methoxyphenyl |
| 15 | A.1 | CH ₃ | 4-methoxyphenyl |
| 16 | A.1 | CH ₃ | 2,4-dimethoxyphenyl |
| 17 | A.1 | CH ₃ CH ₂ | 3,4-dimethoxyphenyl |
| 18 | A.1 | H | 2,4,6-trimethoxyphenyl |
| 9 | A.2 | CH ₃ | 6-dimethylamino-3-pyridinyl |
| 10 | A.2 | CH ₃ | 6-dimethylamino-4-methyl-3-pyridinyl |
| 11 | A.2 | CH ₃ | 6-methyl-5-nitro-2-pyridinyl |
| 12 | A.2 | CH ₃ | 5-chloro-2-pyridinyl |

| Interm. No. | Ex. No. | R ³ | Ar |
|-------------|---------|-----------------|------------------------------|
| 19 | A.1 | CH ₃ | 6-methyl-3-pyridinyl |
| 20 | A.1 | CH ₃ | 3-methyl-5-nitro-2-pyridinyl |

B. Preparation of the final products.

Example B.1

- 5 A mixture of intermediate 2 (21 mg) and *N*-propyl-*N*-cyclopropanemethylamine (75 mg) was heated in a sealed reaction vial at 100°C overnight. Chromatography on a preparative TLC plate with 1:5 ethyl acetate-hexanes gave 3-(2,4-dichlorophenyl)-5-methyl-7-(*N*-propyl-*N*-cyclopropanemethylamino)-pyrazolo[2,3-*a*]pyrimidine (compound 17) (17.4 mg) and 3-(2,4-dichlorophenyl)-5-methyl-7-(*N*-cyclopropane-
- 10 methylamino)-pyrazolo[2,3-*a*]pyrimidine (compound 8) (1 mg). In a similar way, starting from intermediate 2 and (S)-(-)-leucinol respectively (R)-(+)-leucinol, (S)-2[[3-(2,4-dichlorophenyl)-5-methyl-7-pyrazolo[2,3-*a*]pyrimidinyl]amino]-4-methyl-1-pentanol (compound 11) and its R-analog (compound 12) were prepared.

Example B.2

- A solution of intermediate 10 (8 g) and di-*n*-propylamine (13 g) in acetonitrile (50 ml) was heated at reflux for 3 hours. The mixture was filtrated through a short silica gel plug with ethyl acetate and the filtrate was concentrated *in vacuo* to provide a light yellow solid which was recrystallized from ether-hexanes, yielding 8.9 g (93%) of 3-[6-
- 20 (dimethylamino)-4-methyl-3-pyridinyl]-2,5-dimethyl-*N,N*-dipropylpyrazolo[2,3-*a*]pyrimidin-7-amine (compound 53). Compound 53 was also converted to its hydrochloric acid addition salt by dissolving compound 53 (8.1 g) in a mixture of diethyl ether (150 ml) and DCM (50 ml) and treating said mixture with HCl in diethyl ether (1 M, 21.3 ml) dropwise with stirring.
- 25 The resulting off-white solid was collected by filtration, yielding 8.7 g (98%) of 3-[6-(dimethylamino)-4-methyl-3-pyridinyl]-2,5-dimethyl-*N,N*-dipropylpyrazolo[2,3-*a*]pyrimidin-7-amine monohydrochloride.

Example B.3

- Intermediate 3 (15 mg) was dissolved in ethanol (0.5 ml) and stirred in the presence of
- 30 sodium ethoxide (12 mg) for 1 hour. Chromatography on silica gel gave 3-(2,4-dichlorophenyl)-7-(ethoxy)-5-methyl-2-methylthio-pyrazolo[2,3-*a*]pyrimidine (compound 50).

Example B.4

A solution of compound 6 (14 mg) in THF (3 ml) was treated with sodium hydride (60 mg, excess) at room temperature for 5 minutes, followed by iodopropane (0.3 ml). The reaction was stirred at room temperature overnight and quenched with methanol (0.5 ml). The resultant mixture was loaded onto a preparative TLC plate and developed with 1:5 ethyl acetate-hexanes to give 3-(2,4-dichlorophenyl)-5-methyl-7-[*N*-(3-methoxypropyl)-*N*-propylamino]-pyrazolo[2,3-*a*]pyrimidine (compound 45) (5.7 mg) as a colorless oil, 3-(2,4-dichlorophenyl)-5-butyl-7-[*N*-(3-methoxypropyl)-*N*-propylamino]-pyrazolo[2,3-*a*]pyrimidine (compound 46) (4 mg) and 3-(2,4-dichlorophenyl)-5-butyl-7-[3-methoxy-propylamino]-pyrazolo[2,3-*a*]pyrimidine (compound 47) (3.5 mg).

Example B.5

Compound 26 (8 mg) was dissolved in DCM (1 ml) and treated with acetic anhydride (0.1 ml) and pyridine (0.1 ml). The mixture was stirred at room temperature overnight and concentrated in vacuo. The residue was diluted with ethyl acetate and filtrated through a short silica gel column. Concentration of the filtrate gave 3-(2,4-dichlorophenyl)-5-methyl-7-[*N*-(2-acetoxyethyl)-*N*-benzylamino]-pyrazolo[2,3-*a*]pyrimidine (compound 27) (8.6 mg) as a colorless oil.

Example B.6

A solution of compound 40 (20 mg) in DCM (3 ml) was treated with 3-chloroperbenzoic acid (19 mg). The solution was stirred at room temperature for 1 hour. Chromatography on a silica gel plate with ethyl acetate-hexanes (1:1) yielded compound 43 and compound 44.

Example B.7

A mixture of compound 55, palladium on activated carbon (100 mg) in dry ethanol (100 ml) was put on the hydrogenation unit and hydrogenation was carried out at 2.7×10^5 Pa (40 psi) for 2 hours. The reaction mixture was filtered and concentrated. The residue was dissolved in diethyl ether and concentrated, yielding 1.49 g of 3-(5-amino-3-methyl-2-pyridinyl)-2,5-dimethyl-*N,N*-dipropyl-pyrazolo[2,3-*a*]pyrimidine-7-amine (compound 56).

Example B.8

To a stirring solution of compound 56 (1.39 g) and aqueous formaldehyde (6.4 g) in ACN (20 ml) was added sodium cyanoborohydride (743 mg) at 0°C. Glacial acetic

- acid (1 ml) was added and the reaction mixture was stirred at room temperature for 2 hours. The reaction mixture was partitioned between ethyl acetate and saturated aqueous sodium bicarbonate. The organic layer was washed with brine, dried over sodium sulfate, filtered, concentrated to dryness and purified by flash column chromatography on silica gel ($\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$ 150:10:1). The desired fraction was isolated, yielding 1.30 g of 3-[5-(dimethylamino)-3-methyl-2-pyridinyl]-2,5-dimethyl-*N,N*-dipropyl-pyrazolo[2,3-*a*]pyrimidine-7-amine (compound 57).

Example B.9


- To a solution of compound 56 (100 mg) in a mixture of concentrated HCl (1 ml) and water (1 ml), cooled in an ice bath, was added a solution of sodium nitrite (21 mg) in water (1 ml). The solution was added to a mixture of copper(I)chloride (281 mg) in concentrated HCl with stirring in an ice bath. A solid separated and the mixture was heated to 60°C and a clear solution was obtained. The reaction mixture was basified with NaOH, extracted with ethyl acetate, washed with brine and concentrated. Purification by preparative TLC yielded 16 mg of compound 98.




Example B.10

- Compound 56 (100 mg) was added to a solution of fluoroboric acid (48% wt in water, 2 ml), cooled in an ice bath, followed by addition of a solution of sodium nitrite (20 mg) in water (1 ml). The temperature was kept under 10°C. The solid was collected by filtration, dried and partitioned between an aqueous sodium bicarbonate solution and ethyl acetate. The organic layer was washed with brine, concentrated and the residue was purified by preparative TLC yielding 20 mg of compound 100.

Tables 2 to 5 list the compounds that were prepared according to one of the above Examples and tables 6 and 7 list the analytical data for these compounds.


Table 2 :



| Co. No. | Ex. No. | R ⁴ | R ⁵ | Ar |
|---------|---------|------------------|--|--------------------|
| 1 | B.1 | hydrogen | n-propyl | 2,4-dichlorophenyl |
| 2 | B.1 | hydrogen | 2-methylpropyl | 2,4-dichlorophenyl |
| 3 | B.1 | hydrogen | 1,1-dimethylethyl | 2,4-dichlorophenyl |
| 4 | B.1 | hydrogen | 3-hydroxypropyl | 2,4-dichlorophenyl |
| 5 | B.1 | hydrogen | 3-pentyl | 2,4-dichlorophenyl |
| 6 | B.1 | hydrogen | CH ₃ O(CH ₂) ₃ - | 2,4-dichlorophenyl |
| 7 | B.1 | hydrogen | (CH ₃) ₂ CHO(CH ₂) ₃ - | 2,4-dichlorophenyl |
| 8 | B.1 | hydrogen | cyclopropylmethyl | 2,4-dichlorophenyl |
| 9 | B.1 | hydrogen | 3-methyl-2-butyl | 2,4-dichlorophenyl |
| 10 | B.1 | hydrogen | 4-methyl-2-pentyl | 2,4-dichlorophenyl |
| 11 | B.1 | hydrogen |  (S) | 2,4-dichlorophenyl |
| 12 | B.1 | hydrogen |  (R) | 2,4-dichlorophenyl |
| 13 | B.1 | hydrogen |  (S) | 2,4-dichlorophenyl |
| 14 | B.3 | methyl | cyclopropylmethyl | 2,4-dichlorophenyl |
| 15 | B.1 | ethyl | n-butyl | 2,4-dichlorophenyl |
| 16 | B.2 | n-propyl | n-propyl | 2,4-dichlorophenyl |
| 17 | B.1 | n-propyl | cyclopropylmethyl | 2,4-dichlorophenyl |
| 18 | B.1 | n-propyl | HOCH ₂ CH ₂ - | 2,4-dichlorophenyl |
| 19 | B.1 | n-propyl | CH ₃ O(CH ₂) ₂ - | 2,4-dichlorophenyl |
| 20 | B.2 | n-propyl | n-propyl | 4-chlorophenyl |
| 21 | B.1 | n-propyl | cyclopropyl | 2,6-dichlorophenyl |
| 22 | B.1 | n-propyl | CH ₃ O(CH ₂) ₂ | 2,6-dichlorophenyl |
| 23 | B.4 | n-butyl | cyclopropylmethyl | 2,4-dichlorophenyl |
| 24 | B.1 | 2-methylpropyl | cyclopropylmethyl | 2,4-dichlorophenyl |
| 25 | B.1 | 3-methyl-2-butyl | cyclopropylmethyl | 2,4-dichlorophenyl |

| Co. No. | Ex. No. | R ⁴ | R ⁵ | Ar |
|---------|---------|--|--|--|
| 26 | B.1 | HOCH ₂ CH ₂ - | phenylmethyl | 2,4-dichlorophenyl |
| 27 | B.5 | CH ₃ COO(CH ₂) ₂ - | phenylmethyl | 2,4-dichlorophenyl |
| 28 | B.5 | CH ₃ COO(CH ₂) ₂ - | n-propyl | 2,4-dichlorophenyl |
| 29 | B.4 | allyl | cyclopropylmethyl | 2,4-dichlorophenyl |
| 45 | B.4 | n-propyl | CH ₃ O(CH ₂) ₃ | 2,4-dichlorophenyl |
| 58 | B.1 | CH ₃ O(CH ₂) ₂ - | CH ₃ O(CH ₂) ₂ - | 2,4-dichlorophenyl |
| 59 | B.1 | hydrogen | CH ₃ OCH ₂ CH(CH ₃)- | 2,4-dichlorophenyl |
| 60 | B.1 | hydrogen | 1-hydroxy-2-hexyl | 2,4-dichlorophenyl |
| 61 | B.1 | hydrogen | 1-hydroxy-2-pentyl | 2,4-dichlorophenyl |
| 62 | B.2 | n-propyl | n-propyl | 6-dimethylamino- 2,4-dimethyl-3-pyridinyl |
| 63 | B.1 | ethyl | n-butyl | 6-dimethylamino- 2,4-dimethyl-3-pyridinyl |
| 64 | B.1 | n-propyl | cyclopropylmethyl | 6-dimethylamino- 2,4-dimethyl-3-pyridinyl |
| 65 | B.1 | n-propyl | cyclopropylmethyl | 6-methyl-3-pyridinyl |
| 66 | B.1 | n-butyl | n-butyl | 6-methyl-3-pyridinyl |
| 67 | B.1 | n-propyl | cyclopropylmethyl | 2,4-dimethoxyphenyl |
| 68 | B.2 | n-propyl | n-propyl | 2,4,6-trimethoxyphenyl |
| 69 | B.1 | n-propyl | CH ₃ O(CH ₂) ₂ - | 2,4,6-trimethoxyphenyl |
| 70 | B.1 | n-propyl | cyclopropylmethyl | 2,4,6-trimethoxyphenyl |
| 71 | B.1 | ethyl | n-butyl | 2,4,6-trimethoxyphenyl |
| 72 | B.1 | ethyl | ethyl | 2,4,6-trimethoxyphenyl |

Table 3 :




5


| Co. No. | Ex. No. | R ³ | R ⁴ | R ⁵ | Ar |
|---------|---------|-------------------|----------------|----------------|--------------------|
| 30 | B.1 | CH ₃ S | hydrogen | n-propyl | 2,4-dichlorophenyl |
| 31 | B.1 | CH ₃ S | hydrogen | 2-propyl | 2,4-dichlorophenyl |

| Co. No. | Ex. No. | R ³ | R ⁴ | R ⁵ | Ar |
|---------|---------|---------------------------------|----------------|-----------------------|---------------------|
| 32 | B.1 | CH ₃ S | hydrogen | 3-heptyl | 2,4-dichlorophenyl |
| 34 | B.1 | CH ₃ S | hydrogen | 2-methoxyphenylmethyl | 2,4-dichlorophenyl |
| 35 | B.1 | CH ₃ S | methyl | methyl | 2,4-dichlorophenyl |
| 36 | B.1 | CH ₃ S | ethyl | ethyl | 2,4-dichlorophenyl |
| 37 | B.2 | CH ₃ S | n-propyl | n-propyl | 2,4-dichlorophenyl |
| 38 | B.1 | CH ₃ S | n-propyl | cyclopropylmethyl | 2,4-dichlorophenyl |
| 39 | B.1 | CH ₃ S | 2-propyl | 2-propyl | 2,4-dichlorophenyl |
| 40 | B.1 | CH ₃ S | n-butyl | n-butyl | 2,4-dichlorophenyl |
| 41 | B.1 | CH ₃ S | n-butyl | cyclopropylmethyl | 2,4-dichlorophenyl |
| 42 | B.1 | CH ₃ S | allyl | allyl | 2,4-dichlorophenyl |
| 43 | B.6 | CH ₃ SO | n-propyl | cyclopropylmethyl | 2,4-dichlorophenyl |
| 44 | B.6 | CH ₃ SO ₂ | n-propyl | cyclopropylmethyl | 2,4-dichlorophenyl |
| 73 | B.2 | CH ₃ CH ₂ | n-propyl | n-propyl | 3,4-dimethoxyphenyl |

Table 4 :



| Co. No. | Ex. No. | R ⁴ | R ⁵ | Ar |
|---------|---------|--|--|--------------------------------------|
| 48 | B.2 | n-propyl | n-propyl | 4-chlorophenyl |
| 49 | B.1 | n-propyl | cyclopropylmethyl | 2,4-dichlorophenyl |
| 51 | B.1 | n-propyl | CH ₃ O(CH ₂) ₂ - | 2,4-dimethoxyphenyl |
| 52 | B.2 | n-propyl | n-propyl | 6-dimethylamino-3-pyridinyl |
| 53 | B.2 | n-propyl | n-propyl | 6-dimethylamino-4-methyl-3-pyridinyl |
| 54 | B.2 | n-propyl | n-propyl | 5-chloro-2-pyridinyl |
| 55 | B.2 | n-propyl | n-propyl | 3-methyl-5-nitro-2-pyridinyl |
| 56 | B.7 | n-propyl | n-propyl | 5-amino-3-methyl-2-pyridinyl |
| 57 | B.8 | n-propyl | n-propyl | 5-dimethylamino-3-methyl-2-pyridinyl |
| 74 | B.1 | CH ₃ O(CH ₂) ₂ - | CH ₃ O(CH ₂) ₂ - | 4-chlorophenyl |
| 75 | B.1 | HOCH ₂ CH ₂ - | phenylmethyl | 4-chlorophenyl |

| Co. No. | Ex. No. | R ⁴ | R ⁵ | Ar |
|---------|---------|----------------|--|--|
| 76 | B.1 | hydrogen | 1-hydroxy-2-hexyl | 4-chlorophenyl |
| 77 | B.1 | hydrogen | 1-hydroxy-2-pentyl | 4-chlorophenyl |
| 78 | B.1 | hydrogen | CH ₃ S(CH ₂) ₂ - | 4-chlorophenyl |
| 79 | B.1 | hydrogen |  (S) | 4-chlorophenyl |
| 80 | B.1 | hydrogen | 1-hydroxy-2-hexyl | 6-dimethylamino-4-methyl-3-pyridinyl |
| 81 | B.1 | ethyl | n-butyl | 6-dimethylamino-4-methyl-3-pyridinyl |
| 82 | B.1 | n-propyl | cyclopropylmethyl | 6-dimethylamino-4-methyl-3-pyridinyl |
| 83 | B.1 | n-propyl | phenylmethyl | 6-dimethylamino-4-methyl-3-pyridinyl |
| 84 | B.1 | allyl | allyl | 6-dimethylamino-4-methyl-3-pyridinyl |
| 85 | B.1 | n-butyl | n-butyl | 6-dimethylamino-4-methyl-3-pyridinyl |
| 86 | B.1 | hydrogen | 3-pentyl | 6-dimethylamino-4-methyl-3-pyridinyl |
| 87 | B.1 | hydrogen | 2-propyl | 6-dimethylamino-4-methyl-3-pyridinyl |
| 88 | B.1 | hydrogen | 4-methyl-2-pentyl | 6-dimethylamino-4-methyl-3-pyridinyl |
| 89 | B.1 | methyl | n-butyl | 6-dimethylamino-4-methyl-3-pyridinyl |
| 90 | B.2 | n-propyl | n-propyl | 6-dimethylamino-2-methyl-3-pyridinyl |
| 91 | B.1 | n-propyl | cyclopropylmethyl | 6-dimethylamino-2-methyl-3-pyridinyl |
| 92 | B.1 | ethyl | n-butyl | 6-dimethylamino-2-methyl-3-pyridinyl |
| 93 | B.1 | n-butyl | n-butyl | 6-dimethylamino-2-methyl-3-pyridinyl |
| 94 | B.1 | n-propyl | n-propyl | 4,6-dimethyl-3-pyridinyl |
| 95 | B.1 | n-propyl | cyclopropylmethyl | 4,6-dimethyl-3-pyridinyl |
| 96 | B.2 | n-propyl | n-propyl | 6-dimethylamino-2,4-dimethyl-3-pyridinyl |
| 97 | B.1 | n-propyl | cyclopropylmethyl | 6-dimethylamino-2,4-dimethyl-3-pyridinyl |

| Co. No. | Ex. No. | R ⁴ | R ⁵ | Ar |
|---------|---------|--|--|--|
| 98 | B.9 | n-propyl | n-propyl | 5-chloro-3-methyl-2-pyridinyl |
| 99 | B.9 | n-propyl | n-propyl | 5-iodo-3-methyl-2-pyridinyl |
| 100 | B.10 | n-propyl | n-propyl | 5-fluoro-3-methyl-2-pyridinyl |
| 101 | B.2 | n-propyl | n-propyl | 3-chloro-5-trifluoromethyl-2-pyridinyl |
| 102 | B.1 | CH ₃ O(CH ₂) ₂ - | CH ₃ O(CH ₂) ₂ - | 3-chloro-5-trifluoromethyl-2-pyridinyl |
| 103 | B.2 | n-propyl | n-propyl | 3,5-dichloro-2-pyridinyl |
| 104 | B.1 | n-propyl | CH ₃ O(CH ₂) ₂ - | 3,5-dichloro-2-pyridinyl |
| 105 | B.1 | n-propyl | cyclopropylmethyl | 3,5-dichloro-2-pyridinyl |
| 106 | B.2 | n-propyl | n-propyl | 3-methyl-5-methoxy-2-pyridinyl |
| 107 | B.8 | n-propyl | n-propyl | 5-diethylamino-3-methyl-2-pyridinyl |
| 33 | B.8 | n-propyl | n-propyl | 6-diethylamino-4-methyl-3-pyridinyl |
| 108 | B.8 | n-propyl | n-propyl | 5-N-piperidinyl-3-methyl-2-pyridinyl |
| 109 | B.2 | n-propyl | n-propyl | 5-methyl-3-nitro-2-pyridinyl |
| 110 | B.2 | n-propyl | n-propyl | 3-amino-5-methyl-2-pyridinyl |
| 111 | B.9 | n-propyl | n-propyl | 3-chloro-5-methyl-2-pyridinyl |
| 112 | B.2 | n-propyl | n-propyl | 3-methylamino-5-methyl-2-pyridinyl |
| 113 | B.2 | n-propyl | n-propyl | 3-dimethylamino-5-methyl-2-pyridinyl |
| 114 | B.2 | n-propyl | n-propyl | 2-methyl-5-pyridinyl |
| 115 | B.1 | n-propyl | cyclopropylmethyl | 4-isopropylphenyl |
| 116 | B.2 | n-propyl | n-propyl | 3,4-dimethoxyphenyl |
| 117 | B.1 | n-propyl | CH ₃ O(CH ₂) ₂ - | 3,4-dimethoxyphenyl |
| 118 | B.1 | n-propyl | cyclopropyl | 2,4-dimethoxyphenyl |
| 119 | B.1 | hydrogen | CH ₃ O(CH ₂) ₂ - | 2,4-dimethoxyphenyl |
| 120 | B.1 | n-propyl | HO(CH ₂) ₂ - | 2,4-dimethoxyphenyl |
| 121 | B.8 | n-propyl | n-propyl | 5-methylamino-3-methyl-2-pyridinyl |
| 122 | B.2 | n-propyl | n-propyl | 2,4-dimethoxyphenyl |
| 123 | B.1 | ethyl | n-butyl | 2,4-dimethoxyphenyl |
| 124 | B.1 | n-propyl | 2-hydroxypropyl | 2,4-dimethoxyphenyl |
| 125 | B.2 | n-propyl | n-propyl | 4-methoxyphenyl |
| 126 | B.2 | n-propyl | n-propyl | 3-methoxyphenyl |
| 127 | B.1 | n-propyl | cyclopropylmethyl | 3-methoxyphenyl |








| Co. No. | Ex. No. | R ⁴ and R ⁵ taken together | Ar |
|---------|---------|---|--------------------------------------|
| 128 | B.1 |  | 4-chlorophenyl |
| 129 | B.1 |  | 4-chlorophenyl |
| 130 | B.1 |  | 6-dimethylamino-4-methyl-3-pyridinyl |
| 131 | B.1 |  | 6-dimethylamino-4-methyl-3-pyridinyl |

Table 5 :



| Co. No. | Ex. No. | R ¹ | R ² | R ³ | Ar |
|---------|---------|---|---------------------------------|---------------------|--------------------|
| 46 | B.4 |  | n-C ₄ H ₉ | H | 2,4-dichlorophenyl |
| 47 | B.4 |  | n-C ₄ H ₉ | H | 2,4-dichlorophenyl |
| 50 | B.3 | CH ₃ CH ₂ O- | CH ₃ | CH ₃ -S- | 2,4-dichlorophenyl |

5

Table 6 : Analytical data

| Co. No. | ¹ H NMR data (CDCl ₃) | MS M + 1 |
|---------|--|-------------|
| 1 | δ 0.45 (m,2H), 0.78 (m,2H), 1.20 (m,1H), 2.84 (s,3H), 3.40 (d,2H), 5.98 (s,1H), 7.41 (dd,1H), 7.52 (d,1H), 7.76 (d,1H), 8.32 (s,1H) | 335 |
| 2 | δ 0.88 (d,3H), 1.05 (d,3H), 1.74 (m,1H), 2.52 (s,3H), 3.20 (t,2H), 5.87 (s,1H), 6.40 (t,1H), 7.30 (dd,1H), 7.47 (d,1H), 7.99 (d,1H), 8.42 (s,1H) | 349 |

| Co. No. | ¹ H NMR data (CDCl ₃) | MS M + 1 |
|---------|---|-------------|
| 3 | δ 1.57 (s,9H), 2.54 (s,3H), 6.05 (s,1H), 6.48 (s,1H), 7.31 (d,1H), 7.47 (s,1H), 7.99 (d,1H), 8.39 (s,1H) | 349 |
| 4 | δ 1.97 (m,2H), 2.43 (s,3H), 3.62 (m,2H), 3.75 (t,3H), 4.00 (brs, 1H), 6.18 (s,1H), 7.39 (dd,1H), 7.52 (d,1H), 8.19 (d,1H), 8.43 (s,1H) | 351 |
| 5 | δ 1.03 (t,6H), 1.60-1.85 (m,4H), 2.56 (s,3H), 3.49 (m,1H), 5.92 (s,1H), 6.19 (d,1H), 7.35 (dd,1H), 7.50 (dd,1H), 8.01 (d,1H), 8.44 (s,1H) | 363 |
| 6 | δ 2.02 (m,2H), 2.51 (s,3H), 3.40 (s,3H), 3.51 (m,4H), 5.91 (s,1H), 6.71 (t,1H), 7.31 (dd,1H), 7.47 (d,1H), 7.99 (d,1H), 8.42 (s,1H) | 365 |
| 7 | δ 1.23 (d,6H), 2.00 (m,2H), 2.52 (s,1H), 3.51 (dd,2H), 3.59 (m,2H), 5.90 (s,1H), 6.92 (t,1H), 7.31 (dd, 1H), 7.46 (d,1H), 8.00 (d,1H), 8.41 (s,1H) | 393 |
| 8 | δ 0.35 (m,2H), 0.68 (m,2H), 1.25 (m,1H), 2.52 (s,3H), 3.23 (dd,2H), 5.88 (s,1H), 6.40 (t,1H), 7.32 (d,1H), 7.48 (s,1H), 7.99 (s,1H), 8.43 (s,1H) | 347 |
| 9 | δ 1.03 (d,3H), 1.06 (d,3H), 1.31 (d,3H), 1.95 (m,1H), 2.52 (s,3H), 3.54 (m,1H), 5.88 (s,1H), 6.23 (d,1H), 7.32 (d,1H), 7.47 (s,1H), 7.98 (d,1H), 8.41 (s,1H) | 363 |
| 10 | δ 0.95 (d,3H), 0.97 (d,3H), 1.40-1.80 (m,3H), 2.53 (s,3H), 3.76 (m,1H), 5.89 (s,1H), 6.12 (d,1H), 7.32 (d,1H), 7.47 (s,1H), 7.97 (d,1H), 8.40 (s,1H) | 377 |
| 11 | δ 0.94 (d,3H), 0.99 (d,3H), 1.53 (m,1H), 1.76 (m,1H), 2.50 (s,3H), 2.93 (brs,1H), 3.48 (m,1H), 3.76 (m,1H), 5.97 (s,1H), 6.22 (d,1H), 7.35 (dd, 1H), 7.51 (d,1H), 7.95 (d,1H), 8.36 (s, | 393 |
| 12 | δ 0.92 (d,3H), 0.97 (d,3H), 1.52 (m,2H), 1.72 (m,1H), 2.47 (s,3H), 3.02 (brs,1H), 3.42 (m,1H), 3.72 (m,2H), 5.94 (s,1H), 6.20 (d,1H), 7.33 (d,1H), 7.49 (s,1H), 7.92 (d,1H), 8.33 (s, | 393 |
| 13 | δ 0.99 (d,6H), 1.98 (m,1H), 2.44 (s,3H), 3.22 (brs, 1H), 3.46 (m,2H), 3.73 (m,1H), 5.88 (s,1H), 6.31 (d,1H), 7.32 (d,1H), 7.48 (s,1H), 7.92 (d,1H), 8.32 (s, 1H) | 379 |
| 14 | δ 0.21 (m,2H), 0.54 (m,2H), 1.10 (m,1H), 2.53 (s,3H), 3.28 (s,3H), 3.91 (d,2H), 5.99 (s,1H), 7.32 (d,1H), 7.48 (s,1H), 7.97 (d,1H), 8.43 (s,1H) | 379 |

| Co. No. | ¹ H NMR data (CDCl ₃) | MS M + 1 |
|---------|---|-------------|
| 15 | δ 0.96 (t,3H), 1.31 (t,3H), 1.38 (m,2H), 1.71 (m,2H), 2.49 (s,3H), 3.74 (t,2H), 3.65 (q,2H), 5.90 (s,1H), 7.30 (dd,1H), 7.46 (d,1H), 7.97 (d,1H), 8.40 (s,1H) | 377 |
| 16 | δ 0.96 (t,6H), 1.73 (m,4H), 2.49 (s,3H), 3.73 (t,4H), 5.88 (s,1H), 7.33 (dd,1H), 7.46 (d,1H), 7.98 (d,1H), 8.40 (s,1H) | 377 |
| 17 | δ 0.26 (m,2H), 0.58 (m,2H), 0.98 (t,3H), 1.18 (m,1H), 1.76 (m,2H), 2.53 (s,3H), 3.76 (d,2H), 3.81 (t,2H), 6.02 (s,1H), 7.34 (dd,1H), 7.49 (dd,1H), 8.44 (s,1H) | 389 |
| 18 | δ 1.04 (t,3H), 1.77 (m,2H), 2.53 (s,3H), 3.47 (t,2H), 3.98 (m,2H), 4.06 (m,2H), 5.52 (brs, 1H), 6.03 (s,1H), 7.33 (d,1H), 7.48 (s,1H), 7.93 (d, 1H), 8.39 (s, 1H) | 379 |
| 19 | δ 0.98 (t,3H), 1.76 (m,2H), 2.50 (s,3H), 3.35 (s,3H), 3.67 (t,2H), 3.72 (m,2H), 4.14 (m,2H), 5.95 (s,3H), 7.29 (d,1H), 7.47 (s,1H), 7.97 (d,1H), 8.39 (s,1H) | 393 |
| 20 | δ 0.96 (t,6H), 1.73 (m,4H), 2.55 (s,3H), 3.74 (t,4H), 5.90 (s,1H), 7.40 (d,2H), 8.04 (d,2H), 8.25 (s,1H) | 343 |
| 21 | δ 0.26 (m,2H), 0.51 (m,2H), 0.92 (t,3H), 1.10 (m,1H), 1.69 (m,2H), 2.72 (s,3H), 3.41 (d,2H), 3.47 (t,2H), 6.17 (s,1H), 7.18 (t,1H), 7.43 (d,2H), 7.99 (s,1H) | 389 |
| 22 | δ 0.93 (t,3H), 1.67 (m,2H), 2.71 (s,3H), 3.32 (s,3H), 3.44 (t,2H), 3.61 (m,2H), 3.68 (m,2H), 6.16 (s,1H), 7.17 (t,1H), 7.41 (d,2H), 7.99 (s,1H) | 393 |
| 23 | δ 0.24 (m,2H), 0.55 (m,2H), 0.95 (t,3H), 1.38 (m,2H), 1.69 (m,2H), 2.51 (s,3H), 3.74 (d,2H), 3.82 (t,2H), 6.00 (s,1H), 7.31 (dd,1H), 7.48 (dd,1H), 7.98 (d,1H), 8.42 (s,1H) | 403 |
| 24 | δ 0.28 (m,2H), 0.50 (m,2H), 0.94 (d,6H), 1.10 (m,1H), 2.05 (m,1H), 2.52 (s,3H), 3.62 (d,2H), 3.82 (d,2H), 6.03 (s,1H), 7.32 (d,1H), 7.47 (s,1H), 7.99 (d,1H), 8.43 (s,1H) | 403 |
| 25 | δ 0.17 (m,2H), 0.47 (m,2H), 0.97 (t,6H), 1.05 (m,1H), 1.67 (m,4H), 2.52 (s,3H), 2.43 (d,2H), 4.71 (m,1H), 6.13 (s,1H), 7.32 (dd,1H), 7.47 (d,1H), 8.02 (d,1H), 8.43 (s,1H) | 417 |
| 26 | δ 2.41 (s,3H), 4.01 (m,2H), 4.15 (m,2H), 4.79 (s,2H), 5.62 (brs, 1H), 6.02 (s,1H), 7.35 (m,6H), 7.48 (s,1H), 7.91 (d,1H), 8.41 (s,1H) | 427 |
| 27 | δ 1.91 (s,3H), 2.47 (s,3H), 4.20 (t,2H), 4.43 (t,2H), 4.96 (s,2H), 6.03 (s,1H), 7.34 (m,6H), 7.49 (s,1H), 7.94 (d,1H), 8.44 (s,1H) | 469 |

| | | |
|----|--|-----|
| 28 | δ 0.98 (t,3H), 1.74 (m,2H), 1.93 (s,3H), 2.49 (s,3H), 3.59 (m,2H), 4.84 (m,2H), 4.39 (m,2H), 5.95 (s,1H), 7.30 (dd,1H), 7.46 (d,1H), 7.92 (d,1H), 8.38 (s,1H) | 421 |
| 29 | δ 0.28 (m,2H), 0.60 (m,2H), 1.20 (m,1H), 2.51 (s,3H), 3.70 (d,2H), 4.50 (d,2H), 5.29 (d,1H), 5.30 (d,1H), 5.96 (m,1H), 6.05 (s,1H), 7.32 (dd,1H), 7.48 (d,1H), 7.96 (d,1H), 8.44 (s,1H) | 387 |
| 30 | δ 1.07 (t,3H), 1.80 (m,2H), 2.46 (s,3H), 2.55 (s,3H), 3.36 (dt,2H), 5.82 (s,1H), 6.29 (t,1H), 7.29 (d,1H), 7.38 (d,1H), 7.51 (s,1H) | 381 |
| 31 | δ 1.41 (d,6H), 2.46 (s,3H), 2.55 (s,3H), 3.85 (m,1H), 5.82 (s,1H), 6.08 (d,1H), 7.29 (d,1H), 7.39 (d,1H), 7.51 (s,1H) | 381 |
| 32 | δ 0.92 (t,3H), 1.01 (t,3H), 1.25-1.80 (m,8H), 2.45 (s,3H), 2.54 (s,3H), 3.48 (m,1H), 5.79 (s,1H), 6.02 (d,1H), 7.27 (d,1H), 7.39 (d,1H), 7.50 (s,1H) | 437 |
| 33 | δ 0.96 (t,3H), 1.20 (t,6H), 1.74 (m, 4H), 2.15 (s,3H), 2.34 (s,3H), 2.41 (s, 3H), 3.56 (t,4H), 3.79 (t,4H), 5.77 (s,1H), 6.41 (s,1H), 7.54 (s,1H) | - |
| 34 | δ 2.45 (s,3H), 2.56 (s,3H), 3.93 (s,3H), 4.60 (d,2H), 5.90 (s,1H), 6.70 (t,1H), 6.99 (t,2H), 7.27-7.41 (m,4H), 7.52 (s,1H) | 459 |
| 35 | δ 2.44 (s,3H), 2.60 (s,3H), 3.35 (s,6H), 5.00 (brs,1H), 5.84 (s,1H), 7.29 (d,1H), 7.38 (d,1H), 7.51 (s,1H) | 367 |
| 36 | δ 1.36 (s,6H), 2.42 (s,3H), 2.59 (s,3H), 3.80 (m,4H), 5.80 (s,1H), 7.29 (d,1H), 7.38 (d,1H), 7.51 (s,1H) | 395 |
| 37 | δ 0.98 (t,6H), 1.78 (m,4H), 2.40 (s,3H), 2.59 (s,3H), 3.69 (m,4H), 5.74 (s,1H), 7.28 (d,1H), 7.48 (d,1H) | 423 |
| 38 | δ 0.30 (m,2H), 0.62 (m,2H), 0.99 (t,3H), 1.20 (m,1H), 1.81 (m,2H), 2.49 (s,3H), 2.59 (s,3H), 3.65-3.90 (m,4H), 5.87 (s,1H), 7.29 (d,1H), 7.39 (d,1H), 7.51 (s,1H) | 435 |
| 39 | δ 1.05 (d,3H), 1.38 (d,3H), 2.45 (s,3H), 2.56 (s,3H), 3.40 (m,1H), 5.80 (s,1H), 7.30 (d,1H), 7.37 (d,1H), 7.51 (s,1H) | 423 |
| 40 | δ 0.98 (t,6H), 1.40 (m,4H), 1.75 (m,4H), 2.42 (s,3H), 2.59 (s,3H), 3.71 (m,4H), 5.76 (s,1H), 7.29 (d,1H), 7.39 (d,1H), 7.51 (s,1H) | 451 |
| 41 | δ 0.29 (m,2H), 0.61 (m,2H), 0.93 (t,3H), 1.20-1.56 (m,3H), 1.75 (m,2H), 2.43 (s,3H), 2.59 (s,3H), 3.65-3.90 (m,4H), 5.88 (s,1H), 7.30 (d,1H), 7.39 (d,1H), 7.51 (s,1H) | 449 |
| 42 | δ 2.42 (s,3H), 2.59 (s,3H), 4.34 (d,4H), 5.30 (m,4H), 5.87 (s,1H), 6.05 (m,2H), 7.29 (d,1H), 7.39 (d,1H), 7.51 (s,1H) | 419 |

| Co. No. | ¹ H NMR data (CDCl ₃) | MS M + 1 |
|---------|---|-------------------------|
| 43 | δ 0.31 (m,2H), 0.62 (m,2H), 1.00 (t,3H), 1.23 (m,1H), 1.81 (m,1H), 2.49 (s,3H), 3.05 (brs,3H), 3.42-3.95 (m,4H), 6.04 (s,1H), 7.33 (d,1H), 7.50 (s,1H), 7.51 (d,1H) | 451 |
| 44 | δ 0.13 (m,2H), 0.48 (m,2H), 0.90 (t,3H), 1.05 (m,1H), 1.60 (m,2H), 2.64 (s,3H), 3.05 (brs,3H), 3.48 (d,2H), 3.71 (m,2H), 6.04 (s,1H), 7.37 (d,1H), 7.54 (s,1H), 7.60 (d,1H) | 485 (+H ₂ O) |
| 45 | δ 0.95 (t,3H), 1.42 (m,2H), 1.98 (m,2H), 2.49 (s,3H), 3.33 (s,3H), 3.56 (m,2H), 3.75 (m,2H), 5.93 (m,1H), 7.31 (d,1H), 7.46 (s,1H), 7.96 (d,1H), 8.40 (s,1H) | 407 |
| 46 | δ 0.91 (t,3H), 0.95 (t,3H), 1.35-2.05 (m,8H), 2.70 (t,2H), 3.33 (s,3H), 3.40 (m,6H), 5.92 (s,1H), 7.30 (d,1H), 7.46 (s,1H), 8.02 (d,1H), 8.41 (s,1H) | 449 |
| 47 | δ 0.87 (t,3H), 1.25 (m,2H), 1.70 (m,2H), 2.04 (m,2H), 2.68 (m,2H), 3.41 (s,3H), 3.56 (m,4H), 5.86 (s,1H), 6.60 (t,1H), 7.31 (d,1H), 7.46 (s,1H), 8.14 (d,1H), 8.50 (s,1H) | 407 |
| 48 | δ 0.95 (t,6H), 1.71 (m,4H), 2.47 (s,3H), 2.55 (s,3H), 3.71 (t,4H), 5.82 (s,1H), 7.39 (d,2H), 7.70 (d,2H) | 357 |
| 49 | δ 0.24 (m,2H), 0.54 (m,2H), 0.96 (t,3H), 1.12 (m,1H), 1.74 (m,2H), 2.35 (s,3H), 2.44 (s,3H), 3.73 (d,2H), 3.76 (t,2H), 5.93 (s,1H), 7.29 (d,1H), 7.35 (d,1H), 7.51 (s,1H) | 403 |
| 50 | δ 1.64 (t,3H), 2.52 (s,3H), 2.60 (s,3H), 4.56 (q,2H), 6.02 (s,3H), 7.30 (d,1H), 7.37 (d,1H), 7.51 (s,1H) | 368 |
| 52 | δ 0.9 (t, 6H), 1.68 (m, 4H), 2.13 (s, 3H), 2.30 (s, 3H), 2.37 (s, 3H), 3.05 (s, 6H), 3.65 (m, 4H), 5.74 (s, 1H), 6.44 (s, 1H), 8.01 (s, 1H) | M ⁺ = 366 |
| 53 | δ 0.9 (t, 6H), 1.70 (m, 4H), 2.35 (s, 3H), 2.45 (s, 3H), 3.05 (s, 6H), 3.65 (m, 4H), 5.74 (s, 1H), 6.44 (s, 1H), 8.01 (s, 1H) | M ⁺ = 380 |
| 54 | δ 0.95 (t, 6H), 1.73 (m, 4H), 2.52 (s, 3H), 2.76 (s, 3H), 3.73 (t, 4H), 5.85 (s, 1H), 7.68 (dd, 1H), 8.45 (d, 1H), 8.57 (d, 1H) | - |
| 55 | δ 0.96 (t, 6H), 1.70 - 1.78 (m, 4H), 2.44 (s, 3H), 2.47 (s, 3H), 2.50 (s, 3H), 3.73 (t, 4H), 5.48 (s, 1H), 8.36 (d, 1H), 9.32 (d, 1H) | 383 |
| 56 | δ 0.93 (t, 6H), 1.66 - 1.73 (m, 4H), 2.19 (s, 3H), 2.36 (s, 3H), 2.42 (s, 3H), 3.71 (t, 4H), 5.79 (s, 1H), 6.39 (d, 1H), 8.05 (d, 1H) | 353 |
| 57 | δ 0.96 (t, 6H), 1.62 - 1.70 (m, 4H), 2.2 (s, 3H), 2.36 (s, 3H), 2.40 (s, 3H), 2.96 (s, 6H), 3.68 (t, 4H), 5.76 (s, 1H), 6.91 (d, 1H), 8.08 (d, 1H) | 381 |

Table 7 : Analytical data

| Co. No. | Mass spectral data | Co. No. | Mass spectral data |
|---------|------------------------|---------|--|
| 58 | 409 [M ⁺] | 95 | 364 [MH ⁺] |
| 59 | 365 [M ⁺] | 96 | 395 [MH ⁺] |
| 60 | 394 [MH ⁺] | 97 | 407 [MH ⁺] |
| 61 | 380 [MH ⁺] | 98 | 372 [MH ⁺] (Cl ³⁵) |
| 62 | 381 [MH ⁺] | 99 | 464 [MH ⁺] |
| 63 | 381 [MH ⁺] | 100 | 356 [MH ⁺] |
| 64 | 393 [MH ⁺] | 101 | 426 [MH ⁺] (Cl ³⁵) |
| 65 | 350 [MH ⁺] | 102 | 458 [MH ⁺] (Cl ³⁵) |
| 66 | 366 [MH ⁺] | 103 | 392 [MH ⁺] (Cl ³⁵) |
| 67 | 381 [MH ⁺] | 104 | 408 [MH ⁺] (Cl ³⁵) |
| 68 | 399 [MH ⁺] | 105 | 404 [MH ⁺] (Cl ³⁵) |
| 69 | 415 [MH ⁺] | 106 | 368 [MH ⁺] |
| 70 | 411 [MH ⁺] | 107 | 409 [MH ⁺] |
| 71 | 399 [MH ⁺] | 108 | 421 [MH ⁺] |
| 72 | 371 [MH ⁺] | 109 | 383 [MH ⁺] |
| 73 | 397 [MH ⁺] | 110 | 353 [MH ⁺] |
| 74 | 388 [M ⁺] | 111 | 372 [MH ⁺] (Cl ³⁵) |
| 75 | 407 [MH ⁺] | 112 | 366 [MH ⁺] |
| 76 | 373 [MH ⁺] | 113 | 381 [MH ⁺] |
| 77 | 359 [MH ⁺] | 114 | 338 [MH ⁺] |
| 78 | 346 [M ⁺] | 115 | 377 [MH ⁺] |
| 79 | 373 [MH ⁺] | 116 | 383 [MH ⁺] |
| 80 | 397 [MH ⁺] | 117 | 399 [MH ⁺] |
| 81 | 381 [MH ⁺] | 118 | 395 [MH ⁺] |
| 82 | 393 [MH ⁺] | 119 | 415 [MH ⁺] |
| 83 | 429 [MH ⁺] | 120 | 385 [MH ⁺] |
| 84 | 377 [MH ⁺] | 121 | 367 [MH ⁺] |
| 85 | 409 [MH ⁺] | 122 | 383 [MH ⁺] |
| 86 | 367 [MH ⁺] | 123 | 383 [MH ⁺] |
| 87 | 339 [MH ⁺] | 124 | 399 [MH ⁺] |
| 88 | 381 [MH ⁺] | 125 | 338 [MH ⁺] |
| 89 | 367 [MH ⁺] | 126 | 353 [MH ⁺] |
| 90 | 381 [MH ⁺] | 127 | 365 [MH ⁺] |
| 91 | 393 [MH ⁺] | 128 | 370 [M ⁺] |

| Co. No. | Mass spectral data | Co. No. | Mass spectral data |
|---------|------------------------|---------|------------------------|
| 92 | 381 [MH ⁺] | 129 | 368 [M ⁺] |
| 93 | 409 [MH ⁺] | 130 | 393 [MH ⁺] |
| 94 | 352 [MH ⁺] | 131 | 379 [MH ⁺] |

C. Pharmacological examples

Example C.1

REPRESENTATIVE COMPOUNDS HAVING CRF RECEPTOR BINDING

5 ACTIVITY

Compounds were evaluated for binding activity to the CRF receptor by a standard radioligand binding assay as generally described by DeSouza et al. (*J. Neurosci.* 7:88-100, 1987). By utilizing various radiolabeled CRF ligands, the assay may be used
10 to evaluate the binding activity of the compounds of the present invention with any CRF receptor subtype. Briefly, the binding assay involves the displacement of a radiolabeled CRF ligand from the CRF receptor.

More specifically, the binding assay was performed in 1.5 ml Eppendorf tubes using approximately 1×10^6 cells per tube stably transfected with human CRF receptors.
15 Each tube received about 0.1 ml of assay buffer (*e.g.*, Dulbecco's phosphate buffered saline, 10 mM magnesium chloride, 20 μ M bacitracin) with or without unlabeled sauvagine, urotensin I or CRF (final concentration, 1 μ M) to determine nonspecific binding, 0.1 ml of [¹²⁵I] tyrosine - ovine CRF (final concentration ~200 pM or approximately the K_D as determined by Scatchard analysis) and 0.1 ml of a membrane
20 suspension of cells containing the CRF receptor. The mixture was incubated for 2 hours at 22°C followed by the separation of the bound and free radioligand by centrifugation. Following two washes of the pellets, the tubes were cut just above the pellet and monitored in a gamma counter for radioactivity at approximately 80% efficiency. All radioligand binding data was analyzed using a non-linear least-square
25 curve-fitting program.

Binding activity corresponds to the concentration (nM) of the compound necessary to displace 50% of the radiolabeled ligand from the receptor. The following compounds have a $K_i \leq 250$ nM : 5, 10, 12, 15 - 17, 19, 23, 24, 26, 27, 29, 31 - 33, 36 - 43, 48, 49, 51 - 54 and 57 - 130 as listed in Tables 2 - 5. Compounds 17, 29, 38, 41, 42, 48, 49, 51
30 - 54, 57, 58, 70, 71, 81 - 85, 90 - 92, 101, 103 - 105, 115, 118, 121 - 123 and 125 were found to show the best score in this test.

Example C.2CRF STIMULATED ADENYLATE CYCLASE ACTIVITY

The compounds of the present invention may also be evaluated by various functional testing. For example, the compounds of the present invention may be screened for
5 CRF-stimulated adenylate cyclase activity. An assay for the determination of CRF-stimulated adenylate cyclase activity may be performed as generally described by Battaglia et al. (*Synapse* 1:572, 1987), with modifications to adapt the assay to whole cell preparations.

More specifically, the standard assay mixture may contain the following in a final
10 volume of 0.5 ml: 2 mM L-glutamine, 20 mM HEPES, and 1 mM IMBX in DMEM buffer. In stimulation studies, whole cells with the transfected CRF-receptors are plated in 24-well plates and incubated for 1 hour at 37°C with various concentrations of CRF-related and unrelated peptides in order to establish the pharmacological rank-order profile of the particular receptor subtype. Following the incubation, the medium is
15 aspirated, the wells rinsed once gently with fresh medium, and the medium aspirated. To determine the amount of intracellular cAMP, 300 µl of a solution of 95% ethanol and 20 mM aqueous hydrochloric acid is added to each well and the resulting suspensions are incubated at -20°C for 16 to 18 hours. The solution is removed into 1.5
20 ml Eppendorf tubes and the wells washed with an additional 200 µl of ethanol/aqueous hydrochloric acid and pooled with the first fraction. The samples are lyophilized and then resuspended with 500 µl sodium acetate buffer. The measurement of cAMP in the samples is performed using a single antibody kit. For the functional assessment of the compounds, a single concentration of CRF or related peptides causing 80% stimulation
25 of cAMP production is incubated along with various concentrations of competing compounds (10^{-12} to 10^{-6} M).

Example C.3

The plus-maze and defensive withdrawal paradigms are correlated measures of
30 exploratory activity sensitive to anxiogenic and anxiolytic effects of experimental treatments. These animal models are employed to examine anxiolytic and anti-stress actions of compounds of the present invention.

Example C.3-a : The Elevated Plus-Maze Paradigm

This test predicts how animals respond to an approach-avoidance situation involving a
35 bright lighted space versus a dark "safe" area. Both spaces are elevated off the ground and constitute two runways intersecting in the form of plus sign. This type of

approach-avoidance situation is a classical test of "emotionality" and reactivity and is very sensitive to treatments that produce disinhibition (such as sedative/hypnotic drugs) and stress. No motivational constraints are necessary and the animal is free to remain in the dark or venture out on the open arms. The plus-maze apparatus has four arms (10 x 50 cm) at right angles to each other and is elevated from the floor (50 cm). Two of the arms are enclosed with walls (40 cm high) and two arms have no walls (open arms). Subjects are placed individually onto the center of the maze and allowed free access to all four arms for 5 minutes. Subjects are observed through a window in the door and via an on-line display of the rat's location on a computer monitor. Time spent in each arm is recorded automatically by photocell beams and a computer program. The maze is wiped clean with a damp cloth between each trial. The measure of anxiogenic-like behavior in this task is a decrease in time spent on open arms while the measure of anti-stress efficacy is a complementary increase in time spent on open arms.

15 Validation of the Plus-Maze Using CRF Peptides :

Central administration of CRF and exposure to any of several experimental stressors suppresses exploration in the elevated plus maze model of anxiety. When measuring the behavioral response to social defeat, central administration of the alpha-helical CRF (9-41) antagonist either post-stress [25 µg ICV] or pre-stress [1 µg ICV] reverses the anxiogenic effect of social defeat. This anti-stress action of the CFR antagonist is also exerted following intracerebral administration into the central nucleus of amygdala [250 ng IC].

Rats were administered the test compounds orally one hour prior to the five minute test. Some groups were placed in a water-filled pool for ninety seconds immediately prior to placement on the Plus-Maze (Stress group) while control subjects were removed directly from the home cage (Non-Stress group). In the non-treated animal group a significant reduction of the percentage of the average time spent in the open arms was observed (from about 36 to about 16 %). Upon administration of 2.5 or 20 mg/kg of compound 53, the % time in the open arms returned to a level equal, within the error range, to the untreated group.

Example C.3-b : The defensive Withdrawal Paradigm

Testing takes place in a plexiglas open field (106 x 92 x 77 cm) containing a cylindrical galvanized steel chamber measuring 17.1 cm deep and 10.2 cm in diameter. The chamber is open at one end and is located along the wall of the open field aligned

lengthwise and 15.0 cm away from a corner of the open field. The open end faces the corner. The open field is illuminated by fluorescent ceiling lighting. For testing, the animals are introduced into the unfamiliar test environment by placing them into the small chamber. Tests take 5 minutes in duration and the apparatus is cleaned with a mild acetic acid solution after each test. The test compound is administered orally one hour before the 5 minutes test. The behavior of the animals is monitored and recorded by video camera. The latency to leave the chamber will be measured and defined as the placement of all four paws in the open field. Also measured is the number of passages made between the chamber and the open field and the average length of time in the chamber per entry. The measure of anxiolytic efficacy is a decrease in mean time spent within the enclosed chamber. Compound 53 reduced the mean time in the chamber from about 80 seconds to about 20 to 40 seconds when administered at doses of 0.63, 2.5 and 20 mg/kg p.o. to rats.

- 15 Validation of Defensive Withdrawal using CRF peptides
- When injected ICV, both alpha-helical CFR (9-41) and CRF modify behavior in the defensive withdrawal paradigm. In particular, ICV administration of CRF in animals previously familiarized with the apparatus increases both the latency to emerge from the small chamber and mean time spent in the chamber over the fifteen minute session.
- 20 Similarly, infusion of CRF into the locus ceruleus produced similar changes in defensive withdrawal behavior suggesting that the interaction of CRF with noradrenergic neurons is involved in defensive withdrawal behavior in rats. Conversely, ICV administration of alpha-helical CRF (9-14) or d-Phe CRF (12-41), competitive CRF receptor antagonists, reverses the CRF-like effect of a restraint stressor in familiar environment condition and significantly decreases latency to emerge and mean time in chamber measures in animals unfamiliar with the apparatus.

D. Composition examples

30 The following formulations exemplify typical pharmaceutical compositions in dosage unit form suitable for systemic or topical administration to warm-blooded animals in accordance with the present invention.

"Active ingredient" (A.I.) as used throughout these examples relates to a compound of formula (I), a N-oxide form, a pharmaceutically acceptable acid or base addition salt or a stereochemically isomeric form thereof.

Example D.1 : Oral solutions

- 9 g of methyl 4-hydroxybenzoate and 1 g of propyl 4-hydroxybenzoate are dissolved in 4 l of boiling purified water. In 3 l of this solution are dissolved first 10 g of 2,3-dihydroxybutanedioic acid and thereafter 20 g of the A.I. The latter solution is
- 5 combined with the remaining part of the former solution and 12 l of 1,2,3-propanetriol and 3 l of sorbitol 70% solution are added thereto. 40 g of sodium saccharin are dissolved in 0.5 l of water and 2 ml of raspberry and 2 ml of gooseberry essence are added. The latter solution is combined with the former, water is added q.s. to a volume of 20 l providing an oral solution comprising 5 mg of the A.I. per teaspoonful (5 ml).
- 10 The resulting solution is filled in suitable containers.

Example D.2 : Capsules

- 20 g of the A.I., 6 g sodium lauryl sulfate, 56 g starch, 56 g lactose, 0.8 g colloidal silicon dioxide, and 1.2 g magnesium stearate are vigorously stirred together. The
- 15 resulting mixture is subsequently filled into 1000 suitable hardened gelatin capsules, each comprising 20 mg of the A.I..

Example D.3 : Film-coated tabletsPreparation of tablet core

- 20 A mixture of 100 g of the A.I., 570 g lactose and 200 g starch is mixed well and thereafter humidified with a solution of 5 g sodium dodecyl sulfate and 10 g polyvinylpyrrolidone in about 200 ml of water. The wet powder mixture is sieved, dried and sieved again. Then there are added 100 g microcrystalline cellulose and 15 g hydrogenated vegetable oil. The whole is mixed well and compressed into tablets,
- 25 giving 10.000 tablets, each comprising 10 mg of the active ingredient.

Coating

- To a solution of 10 g methyl cellulose in 75 ml of denaturated ethanol there is added a solution of 5 g of ethyl cellulose in 150 ml of dichloromethane. Then there are added 75 ml of dichloromethane and 2.5 ml 1,2,3-propanetriol. 10 g of polyethylene glycol is
- 30 molten and dissolved in 75 ml of dichloromethane. The latter solution is added to the former and then there are added 2.5 g of magnesium octadecanoate, 5 g of polyvinylpyrrolidone and 30 ml of concentrated colour suspension and the whole is homogenated. The tablet cores are coated with the thus obtained mixture in a coating apparatus.

Example D.4 : Injectable solution


1.8 g methyl 4-hydroxybenzoate and 0.2 g propyl 4-hydroxybenzoate were dissolved in about 0.5 l of boiling water for injection. After cooling to about 50°C there were added while stirring 4 g lactic acid, 0.05 g propylene glycol and 4 g of the A.I. The solution
5 was cooled to room temperature and supplemented with water for injection q.s. ad 1 l volume, giving a solution of 4 mg/ml of A.I. The solution was sterilized by filtration and filled in sterile containers.

Example D.5 : Suppositories

10 3 Grams A.I. was dissolved in a solution of 3 grams 2,3-dihydroxybutanedioic acid in 25 ml polyethylene glycol 400. 12 Grams surfactant and 300 grams triglycerides were molten together. The latter mixture was mixed well with the former solution. The thus obtained mixture was poured into moulds at a temperature of 37 to 38°C to form 100 suppositories each containing 30 mg/ml of the A.I.

Claims

1. A compound of formula



- 5 including the stereoisomers and the pharmaceutically acceptable acid addition salt forms thereof, wherein
- R^1 is NR^4R^5 or OR^5 ;
- R^2 is C_{1-6} alkyl, C_{1-6} alkyloxy or C_{1-6} alkylthio;
- R^3 is hydrogen, C_{1-6} alkyl, C_{1-6} alkylsulfonyl, C_{1-6} alkylsulfoxy or C_{1-6} alkylthio;
- 10 R^4 is hydrogen, C_{1-6} alkyl, mono- or di(C_{3-6} cycloalkyl)methyl, C_{3-6} cycloalkyl, C_{3-6} alkenyl, hydroxy C_{1-6} alkyl, C_{1-6} alkylcarbonyloxy C_{1-6} alkyl or C_{1-6} alkyloxy C_{1-6} alkyl;
- R^5 is C_{1-8} alkyl, mono- or di(C_{3-6} cycloalkyl)methyl, Ar^1CH_2 , C_{3-6} alkenyl, C_{1-6} alkyloxy C_{1-6} alkyl, hydroxy C_{1-6} alkyl, thienylmethyl, furanylmethyl,
- 15 C_{1-6} alkylthio C_{1-6} alkyl, morpholinyl, mono- or di(C_{1-6} alkyl)amino C_{1-6} alkyl, di(C_{1-6} alkyl)amino, C_{1-6} alkylcarbonyl C_{1-6} alkyl, C_{1-6} alkyl substituted with imidazolyl; or a radical of formula $-Alk-O-CO-Ar^1$;
- or R^4 and R^5 taken together with the nitrogen atom to which they are attached may form a pyrrolidinyl, piperidinyl, homopiperidinyl or morpholinyl group,
- 20 optionally substituted with C_{1-6} alkyl or C_{1-6} alkyloxy C_{1-6} alkyl; and
- Ar is phenyl; phenyl substituted with 1, 2 or 3 substituents independently selected from halo, C_{1-6} alkyl, trifluoromethyl, hydroxy, cyano, C_{1-6} alkyloxy, benzyloxy, C_{1-6} alkylthio, nitro, amino and mono- or di(C_{1-6} alkyl)amino; pyridinyl; pyridinyl substituted with 1, 2 or 3 substituents independently
- 25 selected from halo, C_{1-6} alkyl, trifluoromethyl, hydroxy, cyano, C_{1-6} alkyloxy, benzyloxy, C_{1-6} alkylthio, nitro, amino, mono- or di(C_{1-6} alkyl)amino and piperidinyl; and wherein said substituted phenyl may optionally be further substituted with one or more halogens;
- Ar^1 is phenyl; phenyl substituted with 1, 2 or 3 substituents each independently
- 30 selected from halo, C_{1-6} alkyl, C_{1-6} alkyloxy, di(C_{1-6} alkyl)amino C_{1-6} alkyl, trifluoromethyl and C_{1-6} alkyl substituted with morpholinyl; or pyridinyl; and
- Alk is C_{1-6} alkanediyl;

with the proviso that 5-methyl-3-phenyl-7-(phenylmethoxy)-pyrazolo[1,5-a]-pyrimidine and 2,5-dimethyl-7-(methylamino)-3-phenyl-pyrazolo[1,5-a]pyrimidine are not included.


- 5 2. A compound according to claim 1 wherein R¹ is OR⁵ and R⁵ is C₁₋₆alkyl; or R¹ is NR⁴R⁵ and R⁴ is hydrogen, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkylcarbonyloxy-C₁₋₆alkyl or C₃₋₆alkenyl, R⁵ is C₁₋₆alkyl, C₃₋₆alkenyl, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxyC₁₋₆alkyl, phenylmethyl or C₃₋₆cycloalkylmethyl; or R⁴ and R⁵ are taken together with the nitrogen atom to which they are attached to form a

10 pyrrolidinyl, piperidinyl, homopiperidinyl or morpholinyl group, optionally substituted with C₁₋₆alkyl or C₁₋₆alkyloxyC₁₋₆alkyl; R² is C₁₋₆alkyl; R³ is hydrogen, C₁₋₆alkyl or C₁₋₆alkylthio; and Ar is a phenyl substituted with 1, 2 or 3 substituents each independently selected from halo, C₁₋₆alkyl or C₁₋₆alkyloxy; or Ar is a pyridinyl substituted with 1, 2 or 3 substituents each independently selected

15 from halo, amino, nitro, trifluoromethyl, mono- or di(C₁₋₆alkyl)amino, piperidinyl or C₁₋₆alkyl.
- 20 3. A compound according to any of claims 1 to 2 wherein R¹ is NR⁴R⁵ and R⁴ is C₂₋₄alkyl, hydroxyC₁₋₂alkyl, C₃₋₄alkenyl or C₁₋₂alkylcarbonyloxyC₂₋₄alkyl; R⁵ is C₂₋₄alkyl, C₃₋₄alkenyl, hydroxyC₂₋₄alkyl or cyclopropylmethyl; R² is C₁₋₂alkyl; R³ is hydrogen, C₁₋₂alkyl or C₁₋₂alkylthio.
- 25 4. A compound according to any of claims 1 to 3 wherein R¹ is NR⁴R⁵ and R⁴ is C₃₋₄alkyl or allyl; R⁵ is C₂₋₄alkyl, allyl or cyclopropylmethyl; R² is methyl; R³ is methyl; and Ar is phenyl substituted in the 3-, 4-, 6-, 2,4- or 2,4,6-positions with halo, C₁₋₆alkyl or C₁₋₆alkyloxy.
- 30 5. A compound according to any of claims 1 to 3 wherein R¹ is NR⁴R⁵ and R⁴ is C₃₋₄alkyl or allyl; R⁵ is C₃₋₄alkyl, allyl or cyclopropylmethyl; R² is methyl; R³ is methyl; and Ar is 3-pyridinyl substituted on the 4- and/or 6-position with methyl or dimethylamino.
- 35 6. A compound according to claim 1 wherein the compound is 3-[6-(dimethylamino)-3-pyridinyl]-2,5-dimethyl-*N,N*-dipropylpyrazolo[2,3-a]pyrimidin-7-amine, or 3-[6-(dimethylamino)-4-methyl-3-pyridinyl]-2,5-dimethyl-*N,N*-dipropylpyrazolo[2,3-a]pyrimidin-7-amine;


the stereochemically isomeric forms, or the pharmaceutically acceptable acid addition salts thereof.

7. A composition comprising a pharmaceutically acceptable carrier, and as active ingredient a therapeutically effective amount of a compound as claimed in any one of claims 1 to 6.
8. A process for preparing a composition as claimed in claim 7 characterized in that a therapeutically effective amount of a compound as claimed in any one of claims 1 to 6 is intimately mixed with a pharmaceutically acceptable carrier.
9. A compound according to any one of claims 1 to 6 for use as a medicine.
10. The use of compounds of formula




including the stereoisomers and the pharmaceutically acceptable acid addition salt forms thereof, wherein R^1 , R^2 , R^3 and Ar are as defined in claim 1, including the compounds 5-methyl-3-phenyl-7-(phenylmethoxy)-pyrazolo[1,5-a]-pyrimidine and 2,5-dimethyl-7-(methylamino)-3-phenyl-pyrazolo[1,5-a]pyrimidine, for the manufacture of a medicament for treating physiological conditions or disorders arising from the hypersecretion of corticotropin-releasing factor (CRF).


11. A compound of formula (II') wherein the radicals R^2 , R^3 and Ar are as defined in any of claims 1 to 5 and radical W' is hydroxy, halo, mesyloxy or tosyloxy; a stereoisomeric form or an acid addition salt form thereof, with the proviso that compounds wherein Ar is unsubstituted phenyl are not included.



12. A process of preparing a compound as claimed in claim 1, characterized by
 a) reacting an intermediate of formula (II) with an intermediate of formula (III) in a reaction-inert solvent,



- b) *O*-alkylating an intermediate of formula (VI) with an intermediate of formula (VII) in a reaction-inert solvent and in the presence of a suitable base, yielding compounds of formula (I-a), defined as compounds of formula (I) wherein R¹ is OR⁵,
- 10




15 wherein in the above reaction schemes the radicals R¹, R², R³, R⁵ and Ar are as defined in claim 1 and W is an appropriate leaving group;

or, if desired, converting compounds of formula (I) into each other following art-known transformation reactions; and further, if desired, converting the compounds of formula (I), into an acid addition salt by treatment with an acid, or conversely, converting the acid addition salt form into the free base by treatment with alkali; and, if desired, preparing stereochemically isomeric forms thereof.

20

13. A process of preparing a compound of formula (II') as claimed in claim 10 characterized by

25 reacting an intermediate of formula (IV) with a β -keto ester (V) in a reaction-inert solvent, thereby yielding compounds of formula (II'-a), defined as compounds of formula (II') wherein W' is hydroxy;



and optionally converting compounds of formula (II'-a) into compounds of formula (II'-b), defined as compounds of formula (II') wherein W' is other than hydroxy;

5 wherein in the above reaction schemes the radicals R², R³ and Ar are as defined in claim 1 and W' is hydroxy, halo, mesyloxy or tosyloxy;

or, if desired, converting compounds of formula (II') into each other following art-known transformation reactions; and further, if desired, converting the compounds of formula (II'), into an acid addition salt by treatment with an acid, or conversely,
 10 converting the acid addition salt form into the free base by treatment with alkali; and, if desired, preparing stereochemically isomeric forms thereof.

INTERNATIONAL SEARCH REPORT

Intern al Application No
PCT/EP 97/00459

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D487/04 A61K31/495

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|--|-----------------------|
| X | JP 61 057 587 A (SHIONOGI AND CO., LTD., JAPAN) 1986 cited in the application & CHEMICAL ABSTRACTS, vol. 105, no. 25, 22 December 1986 Columbus, Ohio, US; abstract no. 226617e, see abstract | 1 |
| X | JP 42 011 753 A (SHIONOGI AND CO., LTD.) 1967 cited in the application see the whole document & CHEMICAL ABSTRACTS, vol. 67, no. 23, 4 December 1967 Columbus, Ohio, US; abstract no. 108663r, see abstract | 1 |

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

15 May 1997

Date of mailing of the international search report

29. 05. 97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

De Jong, B

INTERNATIONAL SEARCH REPORT

Intern. Appl. No.
PCT/EP 97/00459

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|---|-----------------------|
| X | J. MED. CHEM. (1976), 19(4), 512-16 CODEN: JMCMAR, 1976, XP002030876 NOVINSON, THOMAS ET AL: "Novel heterocyclic nitrofurfural hydrazones. In vivo antitrypanosomal activity" see page 516, compounds 7a, 7b --- | 11 |
| X | J. MED. CHEM. (1981), 24(5), 610-13 CODEN: JMCMAR;ISSN: 0022-2623, 1981, XP002030877 SENGA, KEITARO ET AL: "Synthesis and antischistosomal activity of certain pyrazolo[1,5-a]pyrimidines" see Table II, no. 33,35 --- | 11 |
| A | WO 94 13677 A (PFIZER ;CHEN YUHPYNG LIANG (US)) 23 June 1994 cited in the application see claims ----- | 1-13 |

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No
PCT/EP 97/00459

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| JP 61057587 A | 24-03-86 | NONE | |
| JP 42011753 A | | NONE | |
| WO 9413677 A | 23-06-94 | AU 5728194 A | 04-07-94 |
| | | CA 2150709 A | 23-06-94 |
| | | CN 1094048 A,B | 26-10-94 |
| | | CZ 9501586 A | 15-11-95 |
| | | EP 0674642 A | 04-10-95 |
| | | FI 935675 A | 18-06-94 |
| | | HU 70426 A | 30-10-95 |
| | | JP 7509728 T | 26-10-95 |
| | | NO 952399 A | 16-08-95 |
| | | NZ 259114 A | 24-03-97 |
| | | PL 309359 A | 02-10-95 |
| | | ZA 9309405 A | 15-06-95 |